







# JOSEPHO BANKS,

BARONETTO,

REGIÆ SOCIETATIS LONDINENSIS

EXIMIO PRÆSIDI:

PEREGRINATORI CELEBERRIMO,

INGENII ACUMINE, LABORUM PATIENTIA,

PHYSICIS

PER OMNE ÆVUM COMMENDATISSIMO,

SECUNDUM HUNC

NATURÆ VIVARII

FASCICULUM,

D. D. D.

GEORGIUS SHAW,

FREDERICUS P. NODDER.

# SIR JOSEPH BANKS, BARONET.

PRESIDENT OF THE

ROYAL'SOCIETY

OF LONDON:

TO WHOSE UNWEARIED LABOURS,

ENLARGED KNOWLEDGE,

AND LIBERAL PATRONAGE,

THE SCIENCE OF NATURAL HISTORY

IS SO HIGHLY INDEBTED,

THIS SECOND VOLUME OF THE

NATURALIST'S MISCELLANY,

IS RESPECTFULLY INSCRIBED

BY

GEORGE SHAW,

FREDERICK P. NODDER.



#### MUS SALIENS.

CHARACTER GENERICUS.

Dentes primores inferiores subulati.

Lin. Syst. Nat. p. 79.

CHARACTER SPECIFICUS, &c.

MUS cauda elongata floccosa, femoribus longissimis, digito utrinque spurio ad pedes posticos.

CUNICULUS pumilio faliens, cauda anomala longissima.

Briffon Quadr. 103.

JERBOA.

Shaw Itin. 248.


Mira pedum anticorum et posteriorum inæqualitate distinguitur Jerboa: antici enim ita breves sunt ut ad progrediendum minime valeant, postici autem totius corporis longitudini pares sunt. Hinc sit ut pedibus solummodo posterioribus stet et quiescat, anticosque ad nullum alium usum convertat' quam ad attrahendum cibum, culmos nempe cum spicis tritici, aliasque plantas; aut ad sodiendum cubile subterraneum. Plerumque more avis incedit, erecta scilicet cruribus, quæ pilis rarissimis vestiuntur, aviumque cruribus sunt valde

valde fimilia. Cauda illi prælonga est, apice sasciculato.

Tres vel quatuor exstant Jerboæ species, vel saltem constantes, ut dicuntur, varietates, quæque in Pennantis Historia Quadrupedum, genus distinctum constituunt; quas vero Linnæus, sub genere Muris ordinat.

Pernicissimæ velocitatis sunt Jerboæ, et ab hoste saltibus continuatis longissimis adeo rapide ausugiunt, ut equo generoso vectus aliquis vix ægreque illas assequi possit. Frigoris sunt impatientes, et tempore brumali in domunculis subterraneis sepultæ obdormiunt. Frugibus vescuntur. In hoc potissimum dissert species de qua jam agitur a communi specie, (quæ Mus Jaculus Linnæi) quod crura paulo supra pedes digito utrinque spurio, longo, velut calcari armantur. Non tantum calidas regiones, Syriam nempe et Barbariam, sed et Sibiriæ partes orientales, idque frequenter, incolit Jerboa. Moles illi est quasi magni soricis.





N

London Settished Ang " 12 190 by F. S. Nodeler & Als Brever Stever

#### THE

# SIBIRIAN JERBOA.

�\*\*\*\*\*\*\*\*

GENERIC GHARACTER.

Two long cutting teeth in each jaw. Fore Legs very short: hind legs very long.

SPECIFIC CHARACTER, &c.

JERBOA with a pair of spurious toes on each hind foot.

SIBIRIAN JERBOA.

Pennant Quadr. p. 429.

The Jerboa is remarkable for the strange disproportion between the length of the fore legs and the hind ones; the former being so very short as to be of no use to the animal in walking, while the latter are so long as to equal the whole body. In consequence of this peculiar formation, the creature stands and rests on its hind legs only, and makes no other use of the fore legs than to hold its food, and draw down the stems of wheat and other vegetables on which it seeds, and to scratch the ground in order to form its burrows. It has the general actions and attitudes of a bird, and the legs are covered thinly with short hair, and very much resemble those of birds. The tail is very long and terminates in a tust.

Q 2

There

There are three or four distinct species, or at least permanent varieties of Jerboa, and in the History of Quadrupeds of Mr. Pennant they constitute a particular genus; but Linnæus places them as species of the genus Mus.

The Jerboas are animals of the most surprizing swiftnefs, and on the approach of danger immediately fpring forward by fuccessive leaps, so very nimbly, that it is faid to be very difficult for a man well mounted to They are impatient of cold, and reovertake them. main during the winter in a dormant state in their bur-They feed on vegetables. The particular species here represented differs principally from the common Jerboa or Mus Jaculus of LINNÆUS in having the legs furnished at a little distance above the seet with a pair of spurious toes or spurs. It is found not only in the warmer regions of Barbary and Syria, but in the Eastern part of Siberia in considerable plenty. It is about the fize of a large rat.

# PENNATULA PHOSPHOREA.

#### CHARACTER GENERICUS.

Flores Hydræ, ad marginem denticulatum pinnarum.

Stirps libera, subulata, apice pinnata.

Lin. Syst. Nat. p. 1321.

CHARACTER SPECIFICUS, &c.

PENNATULA stirpe carnosa, rachi scabra, pinnis imbricatis.

Lin. Syst. Nat. p. 1322.

PENNATULA PHOSPHOREA.

Ellis A&t. Angl. v. 53. t. 19. f. 1—5.

PENNA MARINA.

Baubin Hift. 3. p. 802.

Pennatulæ genus constituunt Zoophytorum moventium, more piscis natantium. Formæ sunt plerumque perelegantis. Species quam descripsimus est fortasse vulgatissima. Circa littora Britanniæ frequens reperitur, et non raro adhæret piscatorum lineis. Animal constat corpore, seu velut stipite paulum complanato, et inde in ramos laterales numerosos expanso, quorum parietes interni in tubulos multos dividuntur, e quorum apicibus totidem prodeunt velut hydrarum,

hydrarum, seu polyporum capitula, cum suis tentaculis. Totum igitur animal videtur esse hydra ramosa et valde composita, cujus corpus in stipite, seu parte nuda continetur, atque in numerosissimos ramulos utrinque continuatur, singulo ramulo peculiari suo capite instructo.

Mirum hoc animal lucem adeo phosphoream emittit, ut piscatores pisces prope ludentes solo Pennatulæ lumine distincte perspicere possint. Color ejus vivide ruber est, et communis magnitudo ut repræsentat tabula.

a. Ramulus microfcopio auctus.





Lunden Subtiched lug "1" 1790 by To Poloder & Ca V'is Brewer Streets

#### THE

## PHOSPHORIC PENNATULA,

O R

#### SEA-PEN.

#### GENERIC CHARACTER.

Animal free, or locomotive.

Body (generally) expanding into processes on the upper part.

Processes or branches furnished with rows of tubular denticles.

Polype-head proceeding from each tube.

SPECIFIC CHARACTER, &c.

SEA-PEN with fleshy stem, middle part rough, pinnules imbricated.

BRITISH SEA-PEN.

Ellis Zooph. p.61.

#### THE COMMON SEA-PEN.

The Pennatulæ or Sea-Pens form a genus of locomotive Zoophytes, and fwim in the manner of a fish. They are in general of a very beautiful appearance. The species here exhibited is the most common: it is found in considerable plenty on the British coasts, and

is drawn up in numbers adhering to the baits of fishermen's lines. The animal confists of a somewhat
flattened stem or body, which is surnished with an
internal bone, and dilates into an expanded part consisting of a great many pinnæ or lateral branches,
which are divided on their inner edges into a number
of tubular processes, through each of which is protuded
a part of the animal, resembling the head of a hydra or
polype: the whole animal therefore may be considered
as a very compound or ramissed hydra or polype, the
body of which is contained in the naked part or stem,
and from thence ramisses into a vast number of processes,
each furnished with its particular head.

This curious animal emits a very strong phosphoric light, and it is even so luminous that it is no uncommon circumstance for the fishermen to see the fish which happen to be swimming near it, merely by the light of the Pennatula. Its colour is a bright red or crimson, and the general size that of the figure.

a. One of the pinnæ magnified.

#### DIONÆA MUSCIPULA.

**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*** 

CHARACTER GENERICUS.

Perianthium pentaphyllum.

Corolla pentapetala.

Germen depressum, crenatum.

Stigma patens, fimbriatum.

Cl. Decandria Monogynia.

CHARACTER SPECIFICUS.

DIONÆA foliis ciliatis.

Carnivoro generi varia est acquirendæ prædæ ratio. Ab hoc nempe vi corripitur; ab illo venatione satigatur: alterum venenum lethale morsu instat, et id omne otiosum deglutit, cui ausugiendi potestatem ademerit: est etiam quod multiplici utitur dolo, incautaque animalia in casses suos illaqueat. Non tantum animalibus majoribus inest hæc prædandi cupido, sed et minimis; præcipueque insectis, quorum plurima interdum vel in suam speciem graviter sæviunt.

Quod vero dictu omnino mirum videatur, sunt etiam plantæ ad insectorum damnum et perniciem natæ, quæ vel a viscoso foliorum aut slorum quibus incaute insident, humore implicantur, vel illius partis quam irritabilem nuncupant physici, pressura includuntur. Ita se rem habere certissime demonstrat species illa Apocyni, botanicis

botanicis nomine Apocyni androsæmifolii probe cognita: in cujus floribus antheræ quæ profundius intra corollam latent, adeo convergunt, ut fint quafi tegmen nectariis subjacentibus; quorum in medio manet gutta liquoris dulcis et mellei quo potissimum delectantur infecta. Antheræ levissimo tactu irritabiles, multo arcitius inter se conveniunt; quo fit ut musca, seu aliud insectum floribus insidens, et proboscidem ad mel exugendum e fundo corollæ protendens, antheris irritatis et subito convergentibus irretiatur, misereque detenta pereat.


Est tamen multo mirabilius exemplum irritationis hujus vegetabilis in planta quæ in tabula depingitur. Cum enim foliorum superficies facillime exasperetur, et singulum folium ad margines serie spinularum seu vallorum cingatur, tactumque, subito in longitudinem arctissime convergat; necesse est ut quicquid insectum ibi malo fato advenerit, nifi ocissime aufugiat, non fecus ac mus in muscipula capiatur; plerumque etiam firma et valida pressura occidatur. Irritabiliorem reddit hanc plantam plenus et meridianus folis calor.

America Septentrionali innascitur, in uliginosis Carolinæ locis præcipue reperta. In Angliam circiter vigintiquinque abhinc annos illata est, et a celeberrimo Ellisio nomine Dionææ Muscipulæ descripta.

Quo præcipue confilio infita fit ilii mira hæc indoles quam jam tractavimus, philosophos omnes hactenus

effugit, latebitque, ut credo, posteros.

Magnitudo ejus plerumque est ut in tabula depicta: nonnullas tamen ipse aliquando vidi quibus caulis multo procerior. Numerus staminum non semper certus est, et idem.





# DIONÆA MUSCIPULA,

OR

### VENUS'S FLY-TRAP.

<del></del>

GENERIC CHARACTER.

Calyx five-leaved.

Corolla five-petaled.

Germen depressed and crenated.

Stigma spreading and simbriated.

SPECIFIC CHARACTER.

DIONÆA with ciliated leaves.

Various are the methods by which animals of a carnivorous nature obtain their destined prey; some seizing the victim with resistless violence, or hunting it down till it is exhausted with fatigue, or inslicting a deadly poison by their bite, and thus securing it beyond all possibility of escape, and afterwards swallowing it at leisure: others exerting every artifice of insidious ingenuity, and entrapping the unwary animal into their snares. It is not only amongst creatures of the larger kind that this predactious disposition prevails, but even throughout many of the smallest tribes of nature; and particularly amongst insects, of which several kinds might be adduced which occasionally prey even on their own species.

What

What is still more extraordinary, there are not wanting amongst vegetables fome instances, in which the fmaller animals meet their fate by alighting on the flowers or leaves; being either held fast by a viscous exfudation from the furface, or confined by the preffure of the irritable parts of the plant. One of the most curious instances of this kind is in a species of Apocynum, well known to Botanists by the name of Apocynum androsæmifolium. In the flowers of this plant the antheræ, which are fituated pretty deep in the flower, converge, so as to form a shelter for the nectaria lying below them, and in the centre of which is contained a small quantity of that sweet juice so peculiarly attrac-The antheræ are possessed of a very tive to infects. great degree of irritability, and fuddenly converge much closer on being touched; when therefore a fly, or other small insect alights on the flower and inserts its proboscis into the centre, it is suddenly caught by the converging antheræ, and detained in this miferable fituation till it perishes.

But a still more wonderful example of vegetable irritability occurs in the plant represented on the annexed plate. In this plant the surface of the leaves is irritable in the highest degree, and each leaf being surnished round the edge with a series of spiny processes, and converging longitudinally when irritated, with a very considerable degree of pressure, it follows that whatever insect is so unfortunate as to alight on the leaf, is, sunless it be extremely nimble) caught as effectually as a mouse in a trap, and is even generally squeezed to death by the pressure. The plant is most irritable during the warm part of the day, and in sull sunshine.

It is a native of North America, and is principally found in the fwampy parts of Carolina, from whence it was introduced into this kingdom about twenty-five years ago, and was described by the late Mr. Ellis by the title of Dionæa Muscipula.

What particular purpose in the Oeconomy of Nature is answered by the imprisoning power of this extraordinary vegetable, it is extremely difficult, and perhaps impossible to determine.

The general fize of the plant is as represented in the plate, but I have seen the stem much taller than in the specimen here sigured. The number of stamina is not always constant.

•

\*

, p ~ ~



#### BUCEROS RHINOCEROS.

#### CHARACTER GENERICUS.

Rostrum convexum, curvatum, cultratum, magnum, extrorsum serratum; Frontis calvaria nuda, osseo-gibbosa.

Nares pone rostri basin.

Lingua acuta, brevis.

Pedes gressorii.

Lin. Syst. Nat. p. 153.

CHARACTER SPECIFICUS, &c.

BUCEROS cornu mandibulari frontis recurvato.

Lin. Syst. Nat. p. 153.

HYDROCORAX INDICUS.

Briff. Av. 4. p. 571.

Genus Buceros ab aliis omnibus avibus facillime dignoscitur ob rostrum ingens et immodicum; primo sane intuitu, desormitati potius fortuitæ et enormi, quam constanti persectoque Naturæ artificio similius. Cuinam commodo inserviat rostrum hoc inustatum dissicile est dicere; nec de hac re aliquid conjicere possum præter quod jampridem observarunt plurimi physici; posse nempe illud ea de causa dari, ut contra hosses

hostes potentiores, siquando cum illis confligitur, mezlius decertetur.

In plerisque hujus generis speciebus pars illa superioris maxillæ quæ prominet, recta est: at in specie de qua jam loquimur, recurvatur; unde et ipsa avis nomine Rhinocerosis distinguitur. Indiam incolit Orientalem, et in insula Java, nee non in Sumatra, in insulis etiam Philippensibus frequenter eonspicitur. Carnes putridas sertur comedere, visceraque animalium incredibili lætitia deglutire.

Magnitudine Meleagridi Gallopavoni Linnæi non longe est inserior. Colores sunt albi nigrique, ut repræsentat tabula. Rostrum uneias decem est longum; cornuque seu processum super mandibulam superiorem gerit incurvatum, linea longitudinali nigra notatum. Mandibula superior ad basin rubra est, indeque ad apicem albo-slavescens: inferior pallida est, basi nigra.



## GREAT HORNBILL,

O R

## RHINOCEROS-BIRD.

<u></u>

#### GENERIC CHARACTER.

Bill convex, large, and bent, with a protuberance on the upper mandible.

Nostrils finall, and feated behind the bafe of the bill.

Tongue small? sharp?

Feet formed for walking, viz. three toes forward, one backward.

SPECIFIC CHARACTER, &c.

BUCEROS with a recurved mandibular horn.

HORNED INDIAN RAVEN, or RHINOCE-ROS-BIRD.

Will. Ornithol. p. 127.

The genus to which this bird belongs, is above all others distinguished by the extraordinary structure of the beak, which at first view has rather the appearance of some enormous deformity, or irregular monstrosity, than of a natural production. What is the intent of Nature

Nature in the formation of this fingular beak, is not yet clearly investigated; it has been supposed, (and not without a confiderable degree of probability) that the birds of this genus, having some very powerful enemies, with which they may have occasional conflicts, are provided with this immoderate beak as a defensive weapon.

In most of the species, the prominent part on the upper mandible is of a straight form; but in the present species it is turned backwards in a most singular manner; and it is from this circumstance that the bird has obtained its common title of Rhinoceros-bird. It is a native of the East-Indies, and is often sound in Sumatra and Java, and the Philippine Islands. It is said-to feed on carrion, and to be particularly fond of the entrails of animals.

This bird is not much inferior in fize to a turkey, and its colours are as reprefented in the plate; viz. black and white. The beak is about ten inches long, and the horn or curved process on the upper part, is marked by a longitudinal line of black. The upper mandible is red at the base, and of a whitish yellow as it approaches the tip. The lower one is of a pale colour, with a black base.

## ACARÚS AUTUMNALIS.

**◇**\*

#### CHARACTER GENERICUS.

Pedes octo.

Oculi duo, ad latera capitis.

Tentacula duo, articulata, pediformia.

Lin. Syst. Nat. p. 1022.

#### CHARACTER SPECIFICUS.

ACARUS globofo-ovatus ruber, abdomine poftice fetofo.

Omnibus fere nimis notum est molestissimum hoc animalculum, mensibus præcipue Augusti et Septembris. Acaro vulgari multo minus est, cutique assixum unguibus, præsertim tentaculis duobus brevibus supra pedes anteriores sitis, visu dissicillimum est, nec sine laceratione avelli potest. Quamvis celeriter currat acarus autumnalis, tardior tamen est multis ejusdem generis. Cuicunque corporis parti se assixur ibi tumor magnitudine pisi, cum summa prurigine.

Color huic animalculo est vivide ruber, et ope microscopii, pars corporis posterior setis rigidis albisque
vestiri videtur. Rostrum, quod interdum visibile est,
sæpius intra thecam reconditum, tubulatum est. A
summo capite prodeunt parvula duo spicula, utrinque
extrorsum spectantia. Discurrunt hæc insecta super
super superabilia.


vegetabilia, præcipueque fentiuntur ab illis qui intergramina et segetes ambulaverint.

Narrat Dominus White in historia sua de Selborne in agro Hamptoniensi, cretaceas ibi regiones illis adeo insestari, ut cuniculariorum retia rubro colore tingantur, ipsique homines morsu immodico irritati, non raro febricitent.

De hac acari specie silent LINNÆUS et FABRICIUS: cursim et obiter describitur in Bakeri libro secundo de microscopiis qui et siguram addidit; quæ cum ab exsicato insecto delineata sit, veram similitudinem vix satis exprimit.

Qui ipsissimum animalculum commodissime spectare velit, duobus vitri frustulis leviter interpositum in microscopio examinet. Hoc modo membra omnia sacillime explicabuntur, sine laceratione aut injuria.





Landon, Published Sope to of good of To C. Voller Che 1.15 Bener Street.

# AUTUMNAL ACARUS,

## HARVEST-BUG.

GENERIC CHARACTER.

Eight Legs.

Two Eyes, fituated on the fides of the head. Two Tentacula, jointed, and shaped like feet.

SPECIFIC CHARACTER.

RED GLOBOSE-OVATE ACARUS, with the abdomen briftly behind.

This troublesome insect will make itself sufficiently known to most people during the months of August and September: it is much smaller than a common mite, and can but just be perceived upon the skin, to which it adheres by its claws, and particularly by the two short arms or tentacula situated above the upper legs.

It can scarcely be separated from the skin without violence, when once it has fixed itself: its motion when disengaged is considerably quick, though by no means equal to that of some other species of acari. On the part where it fixes, it causes a tumor, generally about the size of a pea; sometimes much larger, accompanied with a severe itching.

The

The colour of this dimunitive infect is a bright red; and when microscopically examined, the lower part of the body appears to be coated with stiff white brissles. It feems to be provided with a tubular snout, which is generally conceased or sheathed, but which may sometimes be distinctly seen. On the top of the head are two little processes or sharp implements which turn outwards each way. These insects abound on vegetables, and are generally contracted by walking in gardens, amongst long grass, or corn fields.

According to MR. WHITE in his History of Selborne, they abound to an uncommon degree in the chalky diftricts of that part of Hampshire. He relates that he has been affured that the warreners in the chalky downs, are so much infested by them, and that they swarm to so infinite a degree as to discolour their nets, and give them a reddish cast, whilst the men are so bitten as to be thrown into severs. Vid. Hist. Selb. p. 89.

It is a species which seem to have escaped the notice of systematic Naturalists; and is not to be sound either in Linnæus or Fabricius. A slight general description of it is given in Baker's Employment for the Microscope, accompanied by a figure; but as the sigure was evidently taken from a dried specimen, it gives but a very impersect idea of the animal.

The only way of viewing it to advantage is to place it on a piece of glass and to lay another piece of glass immediately upon it, by which means it will be just so much compressed as to expand all its limbs without being injured; in this situation it may be viewed, especially by a compound microscope, to the greatest possible advantage.

## MANTIS GIGAS.

<del></del>

#### CHARACTER GENERICUS.

Caput nutans, maxillosum, palpis instructum. Antennæ setaceæ.

Alæ quatuor, membranaceæ, convolutæ: inferiores plicatæ.

Pedes antici compressi, subtus serrato-denticulati, armati ungue solitario et digito setaceo laterali articulato; postici quatuor, læves, gressorii.

Thorax linearis, elongatus, augustatus.

Lin. Syft. Nat. p. 689.

#### CHARACTER SPECIFICUS.

MANTIS thorace teretiusculo, elytris brevissimis, pedibus spinosis.

Lin. Syst. Nat. p. 689.


plenum,


Formam magis miram et singularem quam quæ huic insecto est, vix animo possumus concipere. Quod si physicos solos audiremus, nec oculis nostris sides esset, dubitari etiam sortasse posset an tale ullud unquam revera extiterit. In insectis hujus classis, larva, seu animal impersectum, ab imagine ipsa, seu insecto adulto non multum abludit, nisi quod alis careat, quarum illi (ut physicorum more loquar) rudimenta tantum sunt. In musæis rarissime conspicitur insectum hoc alatum,

plenum, et perfectum; larvæ quippe folummodo in Europam advehi folitæ funt, in queis alarum, ut jam diximus, rudimenta vix ac ne vix confpici possunt. Istiusmodi larva communiter bacilli ambulantis nomine cognoscitur.

Singulare aliquid huic infecto est, quodque perpaucis aliis sui generis contingit, alas nempe non tantum thecis externis, sed et elytris quasi secundariis muniri, quæ tamen connexa videntur, quæque ipsas sortasse alas ab injuriis securiores reddant.

Mirum et rarum hoc insectum Insulam Amboynam inhabitat; ipsumque specimen unde depingitur hæc nostra sigura, in Musæo Leveriano tam belle exsiccatum, ut nihil sit persectius, asservatur.





London Sublished Sept VI 1790 by F. J. Solder 42. 1. 15 Brewer Street.

## GREAT MANTIS.

#### GENERIC CHARACTER.

Head unfleady: Mouth armed with jaws, and furnished with palpi.

Antennæ setaceous.

Wings four, membranaceous, convoluted: the lower ones plicated.

Feet anterior compressed, servated beneath, armed with a solitary claw and lateral jointed process; posterior sour, smooth, formed for walking.

Thorax linear, elongated, and narrowed.

#### SPECIFIC CHARACTER.

MANTIS with roundish rough thorax, very short elytra, and spiny seet.

Imagination can hardly figure to itself a creature of a more singular appearance than this insect; and had we only received the accounts of authors, without having seen the animal itself, we might be inclined to question the truth of its existence. In insects of this tribe the animal in its incomplete state, or that which is analogous to the caterpillar-state in the butterslytribe, differs not much from the appearance of the in-

fect in its complete form, except in not being furnished with wings.

This creature is very rarely met with in collections in its complete or winged form; being generally feen in the lefs advanced growth before-mentioned, in which the rudiments of the wings are but just visible. In that imperfect state it has commonly been described under the title of the walking stick.

A most singular circumstance in this insect, (and which takes place in but very sew others of the genus) is, that, exclusive of the elytra, or wing cases, there is an additional pair, which may be supposed to serve as a farther guard in securing the wings themselves: this secondary pair however seem to be connate with the wings themselves, so as to make a part of them.

This most curious and uncommon insect is a native of the Island of Amboyna, and the specimen from which the sigure was taken, is now in the highest prefervation in the Leverian Museum.

## PICUS MINIMUS.

## CHARACTER GENERICUS.

Rostrum polyedrum, rectum: apiec cuneato.

Nares pennis setaceis recumbentibus obtectæ.

Lingua teres, lumbriciformis, longissima, mucronata, apice retrorsum aculeata setis.

Pedes scansorii.

Lin. Syft. Nat. p. 173.

## CHARACTER SPECIFICUS.

PICUS griseus, vertice coccineo, occipite nigro albo punctato.

Species Picorum plurimæ exstant; sed omnium specierum hactenus cognitarum hæc sacile minima. Tabula sistit aviculam magnitudine naturali. Cayanam in America Australi incolit.

## LEAST WOODPECKER.

## GENERIC, CHARACTER.

Bill angular, strait, cuneated at the tip. Nostrils covered with reflected bristly feathers.

Tongue cylindric, worm-shaped, very long, sharppointed, and (generally) aculeated at the tip with reflex bristles.

Feet formed for climbing, viz. two toes forward and two backward.

Lin. Syft. Nat.

## SPECIFIC CHARACTER, &c.

CHESNUT-GREY WOODPECKER, with the top of the head crimfon, and the back part black speckled with white.


Le tres petit PIC DE CAYENNE.


Buff. Oif. 7. p. 37.

### MINUTE WOODPECKER.

Lath Synops. vol. 1. p. 596.

The species of Woodpeckers are very numerous; and of all the species yet discovered, this is by far the least. The plate represents it in its natural size. It is a native of Cayenne in South America.





## LACERTA SALAMANDRA.

CHARACTER GENERICUS.

Corpus tetrapodum, ecaudatum, nudum.

Lin. Syft. Nat. p. 359.

CHARACTER SPECIFICUS, &c.

LACERTA cauda tereti brevi, pedibus muticis, palmis tetradactylis, corpore porofo nudo.

Lin. Syft. Nat. p. 371.

SALAMANDRA.

Gefn. Quadr. 80.

SALAMANDRA TERRESTRIS.

Ray Quadr. 273.

Novum omne et inauditum avida femper aure bibit gens humana. Quasi vero ipsa Natura sua non mille habeat miracula, nunquam desuere qui de monstris et prodigiis, reliquaque insani capitis farragine lubentissime scripserint.

Istiusmodi sunt dracones illi ingentes ab antiquis descripti; nec non Lamia, sera nempe quæ corpus Pantheræ squamis vestitum, cum capite et mammis mulieris sortita est. Huc etiam reserenda est Salamandra, quam vulgus inepte putat in igne illæsam vivere.

Est profecto Salamandra parva lacertæ species, in Germania,

Germania, allifque Europæ regionibus minime raga; coloris nigri, apud latera ventremque maculis magnis longifque vivide flavis, five aureis decorati. Maculæ hæ interdum magnitudine et figura variant. Tegitur lime lacerta mentibus, five foraminibus parvis, humorem guo plerumque madet animal, exudantibus. Hi meatus prope caput magis conspicui sunt quam in reliquo corpore. Cauda ad teretem accedit formam, et longitudine est mediocri, gradațim attenuata.

Hoc est illud animal de quo tot miræ et ridiculædivulgatie funt fabulæ, quodque etiam nunc temporis a vulgo, peculiari quadam et ignoța vi ignis ardorem repellere creditur. Est sane nihil aliud hæc vis antipyretica, quam frigidi animalis humida et naturalis temperies, cujus ope paulo longius ab ignis impetu-manet inconfumptum, quam aliarum fubitantiarum major

ficcitas.





## SALAMANDER.

GENERIC CHARACTER.

Body four-footed, tailed, naked.

SPECIFIC CHARACTER.

LIZARD with cylindric fhortish tail and unarmed feet; the fore-feet divided into four toes, the body porous and naked.

There feems to prevail in the human mind an inherent propenfity to the marvellous. Not contented with the variety of real miracles which Nature through all her kingdoms fo liberally exhibits, the world has long been entertained with the histories of ideal wonders and imaginary monsters, which never existed but in the brains of their first describers.

Amongst beings of this class must be reckoned the large dragons described by the old writers; the Lamia, described as having the head and breasts of a woman, with a body like that of a panther, and covered with scales: to this order must also be referred the ideal Salamander of the vulgar; which is thought capable of living unhurt in the sire,

The real Salamander is nothing more than a fmallish lizard, which is found very frequently in Germany, and

and many other parts of Europe. It is of a black colour, ornamented on the fides and belly with large and longish marks or spots of a rich yellow or gold-colour, and which are frequently somewhat irregular in their shape and disposition. It is covered over with pores or small foramina, through which exsudes the moisture with which it is generally covered: these pores are most conspicuous near the head: the tail is of a moderate length, and is roundish, or somewhat cylindrical, and gradually tapers to the end.

Such is the animal of which fo many incredible tales have been recited, and which still continues to be regarded by the ignorant as possessed of the power of repelling the effect of fire; a power which it possesses in no higher degree than a frog, a snail, or any other moist substance, which is not so immediately consumed as those of a drier nature.

## JULUS MAXIMUS.

**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*** 

#### CHARACTER GENERICUS.

Pedes numerosi, duplo utrinque plures quam corporis segmenta.

Antennæ moniliformes.

Palpi duo articulati.

Corpus semicylindricum.

Lin. Syst. Nat. p. 1064.

CHARACTER SPECIFICUS.


JULUS pedibus utrinque 134.


Lin. Syst. Nat. p. 1066.

Hujus generis infecta scolopendris sunt valde assinia; nisi quod corpora habent cylindracea, non complanata; quodque singulis corporis articulis pedes utrinque duo sint, quorum numerus duplo major est quam corporis articuli, cum in Scolopendris sit par.

Animalia hæc, fi aliquo modo perturbantur, in spiram planam se solent contorquere. Os habent maxillis validissimis munitum, quibus morsum immitem non tamen venenosum possunt insligere. Species hic depicta est sui generis maxima. In America invenitur, sylvas et loca obscura perreptans. Species etiam valde

valde similis, si non eadem, in India Orientali exstat. Color his infectis est lucide suscess, paululum nigrans; crura autem, cum corporis parte inferiore subalbicant.





Sunday Lattished Corestings by File toldlor of for A to Brown Son

17

## GREAT JÜLÜS

#### GENERIC CHARACTER:

Feet numerous: twice as many on each fide as the fegments of the body.

Antennæ moniliform.

Palpi (or feelers) two, jointed.

Body femicylindric.

SPECIFIC CHARACTER.

JÜLUS with about 134 feet on each fide.

These infects are very nearly allied to the Scolopendræ or centipedes, but their body, instead of being flattened, as in those creatures, is nearly cylindrical; and every joint of the body is furnished on each side with two pair of seet; so that the number of seet on each side is double the number of joints, whereas in the Scolopendra they are equal.

These animals, when disturbed, roll themselves up in a flat spiral. Their mouth is armed with a very strong pair of jaws, with which they are capable of inflicting a severe bite: they are not however of a poisonous nature. The species here sigured is the largest of the genus. It is sound in America, where it inhabits

U

woods and retired places; and a species extremely refembling it, if not in reality the same, is found in the East-Indies. The colour of the insect is a blackish brown, of a shining surface: the legs are whitish, as is also the under part of the body.





## STRIX NYCTEA.

VAR. STRIATA.

**�**^^^^^^^

CHARACTER GENERICUS.

Rostrum aduncum (absque cera.)

Nares pennis setaceis recumbentibus obtectæ.

Caput grande: auribus oculisque magnis.

Lingua bisida.

Lin. Syst. Nat. p. 131.


### CHARACTER SPECIFICUS.

STRIX capite lævi, corpore albido maculis lunatis diftantibus fufcis.

I in. Syst. Nat. p. 132.

Avis hæc inter varietates est rarissimas ac elegantissimas magnæ et niveæ strigis Americam simul ac Europam septentrionalem incolentis: 'plerumque est tota nivea, maculis aliquot lunatis susco-nigricantibus interspersis: varietas autem de qua jam loquimur, est per totum sere corpus notis numerosissimis suscis obducta, quæ versus collum sensim angustiores siunt, et in breves et interruptas sascias concurrunt, quibus etiam tota avis subtus notata est: caput omnino albet; ut et pedes, qui, ut in aliis omnibus hujus speciei exemplis,

emplis, ad ipfos ungues plumis velleri fimillibus den, fissime vestiuntur: rostrum unguesque nigricant; irides slavent.





#### THE

## SNOWY OWL.

A STRIPED VARIETY.

**◇**\*\*\*\*\*\*\*

GENERIC CHARACTER.

Bill crooked (not furnished with a cere).

Nostrils covered with recumbent bristly feathers.

Head large: Ears and Eyes large.

Tongue bisid.

SPECIFIC CHARACTER, &c.

HORNLESS WHITE OWL, marked with diffant lunated fpots.

SNOWY OWL.

Lath. 1. p. 132.

GREAT WHITE OWL.

Edwards 2. pl. 61.

The bird here figured is a most curious and beautiful variety of the great Snow-white Owl; a species which is a native both of the northern parts of Europe and of America, and in size is equal to the great Eagle-Owl. In its common or general state it is of a snowy whiteness, with here and there a few small scattered crescent-shaped spots of blackish brown; but the specimen now represented is spotted all over the upper surface with numerous marks of brown,

brown, which as they approach the neck become narrower, and run into short interrupted fasciæ or bands, with which also the whole under surface of the bird is ornamented: the head is white; the feet also are perfectly white; and like all other individuals of this species, are most thickly coated with feathers to the very claws, so as to have the appearance of a very deep fur: the bill and claws are black: the irides yellow.

## SIMIA SPHINX?

<u></u>

#### CHARACTER GENERICUS.

Dentes Primores utrinque quatuor, approximati. Laniarii folitarii, longiores, hinc remoti. Molares obtusi.

Lin. Syft. Nat. p. 34. Cl. Mammalia.—Ord. Primates.

#### CHARACTER SPECIFICUS.

SIMIA SEMICAUDATA, ore vibriffato, unguibus acuminatis, natibus calvis?


Lin. Syft. Nat. p. 35.

Simiarum quæ communiter Papiones nominantur, duæ funt species adeo inter se similes, ut revera licet diversæ, plerumque tamen ab auctoribus, qui suis oculis visas sibi invicem collatas non sedulo examinaverint, pro una eademque specie descriptæ sint: Simia nempe Sphinn et Simia Maimon LINNÆI.

Præcipue distinguuntur diversa magnitudine: Sphinge enim multo minor est Maimon: sed discrepat et vultus color: Sphinx enim nasum habet medium sanguinei ruboris, Maimon obscure tantum incarnatum. Sphingis etiam natibus multo vividior

est rubor, regionique lumborum color omnino cærused violaceus, qui in iisdem partibus alterius vix et ne vix conspici potest. Ad cætera vero adeo sunt affines hæ duæ species ut non mirum sit si incertum discrimen plurimos primo visu sefellerit.

Rarissimus est Simiæ Sphingis in Europa conspectus; Maimonis frequentior, et sere quotidianus. Utræque Africæ calidiores regiones inhabitant. Speciatim notandum est nomina et synonima harum specierum scriptores communiter permiscuisse: Linnær enim Maimon multo melius cum hac nostra specie convenit quam cum illa quam ipse nomine Sphingis descripsit. Suspicor etiam (ut plane rem consitear,) hanc ipsam speciem hic depictam non esse Sphingem Linnær. Ad altitudinem quinque pedum et ultra crescit Sphinx; Maimon vix tres superat.







#### THE

## VARIEGATED BABOON.

#### GENERIC CHARACTER.

Front-Teeth in each jaw 4, placed near together. Canine-Teeth folitary, longer than the others, distant from the remaining teeth, or grinders. Grinders obtuse.

### SPECIFIC CHARACTER, &c.

SHORT-TAILED WHISKERED SIMIA, bare behind, with pointed nails, and violet-coloured loins.

#### GREAT BAROON,

Pennant Hist. Quadr. p. 173.

Amongst the species of Simiæ called Baboons there are two, which though really very distinct from each other, have yet so great a general resemblance, that some authors, not having had opportunities of examining and comparing both species, have consounded them with each other; these two animals are the Simia Sphinx and the Simia Maimon of Linnæus.

The most striking difference is in point of size; the Sphinx being by much the largest of the two; but there is also a considerable difference in the colour of

the face: the middle of the nose, which in the Maimon is merely of a dull flesh-colour, is in the Sphinx of a fanguine red: the hinder part of the Sphinx is also of a much more intense red than in the Maimon; and the skin round the region of the loins is tinged with a very strong and sine violet-colour, which is scarce, if at all perceptible in the Maimon. In other particulars, they so much agree that it is no wonder they should have been frequently mistaken for one and the same species.

The S. Sphinx is a much rarer animal than the Maimon, and indeed is but very feldom to be feen in Europe; whereas the Maimon is not uncommon in most exhibitions of animals. Both species are natives of the hotter parts of Africa. It should be particularly observed that a general confusion seems to take place in the descriptions and synonyms of authors relative to these two animals; for the description given by Linneus of his S. Maimon agrees much better with our supposed Sphinx than with the species so named by Linneus. I am even inclined to suspect after all, that the S. Sphinx of Linneus cannot be the animal here represented. The Sphinx grows to the height of upwards of 5 feet: the Maimon rarely exceeds 2 or 3 at farthest.

## PAPILIO MENELAUS.

#### CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavatocapitatæ.

Alæ fedentis erectæ furfumque conniventes (vo-latu diurno).

Lin. Syst. Nat, p. 744.

CHARACTER SPECIFICUS, &c.

PAPILIO alis dentatis; supra cæruleis nitidissimis; subtus nebulosis, punctis suscis.

Lin. Syst. Nat. p. 749. Eq. Achivi.

Merian Surin. t. 53. Cram. Inf. 2. t. 21. Clerk. Icon. t. 21. f. 1.


En superbientis Naturæ miraculum! eximiæ venustatis insectum! vividos adeo colores sortitum, ut vix ulla arte ad vivum possint exprimi.


"Color alarum cæruleus paginæ superioris (inquit Linnæus) adeo politus nitidusque, ut vix simile in rerum Natura conspiciatur." Mirisicus hic paginæ superioris splendor a sobrio et modesto paginæ inserioris colore longissime distat; qui tamen ocellis serrugineis margine nigro circumscriptis, pupillaque albicante Y gemmatus,

gemmatus, sui generis pulchritudinis exemplar non mediocre censetur.

Externa hæc et interna diversitas, (si causam liceat quærere) Papilionem ab avibus sortasse seçuriorem reddat, ut nempe ab hostibus minus facile discerniqueat, dum alis clausis quiescit, quam cum alis expansis pleno splendore effulserit.

Americæ est incola, et ab eruca magna slavescente, spinis nigricantibus armata, originem ducit,





Louden . Intilished . Anderstyges by A.B. Soulder & J. B. S. it wowen Most.

# MENELAUS,

#### OR THE

# SILVER-BLUE BUTTERFLY.

#### GENERIC CHARACTER.

The Antennæ or Horns thickening towards the upper part, and generally terminating in a knob, or club-shaped tip.

The Wings (when fitting) erect, and meeting upwards. (Flight diurnal.)

## SPECIFIC CHARACTER, &c.

BUTTERFLY with indented wings; the upper furface of a brilliant blue, the lower clouded-brown; and marked with large ocellated fpots.

Mer. Surin. t. 53.

Cram. Inf. 2. t. 21. Clerk. ic. t. 21. f. 1.

So uncommonly bright and brilliant is this superbinsect, that it can but faintly be expressed by the utmost efforts of artificial colouring; and may serve as an instance, amongst many others, of the inimitable beauty which Nature alone can produce.

LINNÆUS in his description of this insect observes, that the blue on the upper surface is so polished and lively that scarce any other natural object can come in competition with it. On the contrary the under surface of the same animal exhibits an example of a species of beauty resulting from a varied combination of the plainest and most sober colours; the ground colour being brown slightly streaked with higher shades, and marked by several very large ocellated ferruginous spots with dark rings and white pupils.

If it were not almost bordering on temerity to attempt a reason for this striking difference between the two surfaces of the same insect, one might suppose that this sobriety of colouring on the lower side, is intended in some measure to secure the animal when sitting at rest, with its wings closed, from the depredations of birds, which are less likely to be attracted in this state than by the sull lustre of its expanded plumage.

It is a native of South America, and proceeds from a large yellow caterpillar, befet with black fpines.

## PSITTACUS MAGNIFICUS.

#### CHARACTER GENERICUS.

Rostrum aduncum: mandibula superiore mobili, cera instructa.

Nares in rostri basi.

Lingua carnofa, obtufa, integra.

Pedes scansorii.

Lin. Syst. Nat. p. 139.

#### CHARACTER SPECIFICUS.

PSITTACUS brachyurus fubcristatus niger, fronte humerisque gilvo maculatis, cauda medio rubra nigro fasciata.

Novæ Hollandiæ infula, dignior fane quæ Continens Auftralis vocetur, alter quafi physicorum orbis, plurima animalia, reliquis regionibus incognita pergit fuppeditare. Nulli tamen alii generi uberior accessit specierum copia quam psittacino, quarum lautissimæ vel ipsius Novæ Hollandiæ vel insularum adjacentium sunt incolæ.

Inter species hasce novas vix venustior est Psittaco Magnisico, qui sane mensura maximos Psittacos Maccatos dictos, physicisque notissimos æquat; in illa autem generis divisione recte disponitur, cui cauda non lanceæ ad instar producta et acuminata est, sed apice quadrato, seu ex rectricibus sere æqualibus constat.

Z

Color generalis huic avi niger est. Caput leviter cristatum plumis densissime vestitur, quarum extremitates maculis gilvis, ut plurimum scatere solent. Maculantur quoque humeri eodem-modo, et in aliquibus speciminibus latera corporis semoraque sasciis gilvis anguste transverseque notantur. Rostrum maximum et validissimum cornei est coloris, seu obscure susci. Crura pedesque crassa et robusta, rostroque sere concoloria. Caudæ imo colori, coccineo nempe, transverse superadditæ sunt sascii numerosæ nigerrimæ; et utrinque prope rachin pennarum coccineus color sere in slavum languescit. Basin apicemque cauda penitus nigra est.

Tota avis tam magnificum oftendit aspectum ut inter principes sui generis merito numeretur. Specimina aliquando paululum variant: color scilicet caudæ coccineus non in sascias numerosas nigro distinctas disponitur, sed latam transversamque aream intemerati ruboris in medio exprimit.





# T H E MAGNIFICENT COCKATOO.

GENERIC CHARACTER.

Bill hooked. Upper mandible moveable.

Nostrils round, placed in the base of the bill.

Tongue sleshy, broad, blunt at the end.

Legs short. Toes formed for climbing, viz. two forward and two backward.

SPECIFIC CHARACTER, &c.

EVEN-TAILED BLACK PARROT, with the head fomewhat crefted, the front and fhoulders fpotted with buff, the middle of the tail red croffed with black bars.

THE BANKSIAN COCKATOO.

Latham's Synopf. Suppl. p. 63.

New Holland, which may not unjustly lay claim to the more dignified title of the Southern Continent, may be confidered as a kind of new world to the Naturalist; and has already afforded several animals unknown to every other part of the globe. To no other genus, however, have such large accessions of new species been added as to that of Psittacus; of which some of the most superb kinds appear to be natives of New Holland, and some of the Southern Islands.

 $Z_2$ 

Of

Of these newly-discovered birds, one of the most august in its appearance is that represented on the plate annexed. In fize it is equal to the great Maccaws already so well known to Naturalists; but it belongs to a different section in the genus, and instead of being furnished with a lanceolate tail, as in those birds, it has that part even at the end, or confishing of seathers of nearly equal length.

. The general color of this bird is a deep black: the head is very full of feathers, and flightly crefted, and is commonly marked in front with feveral buff-coloured fpots, owing to the tips of the feathers on that part being of this colour. Several spots of buff are also scattered over the shoulders, and in some specimens the fides and thighs are flightly barred with narrow fasciæ The bill is of a deep brown, or horn-color, and is uncommonly large and ftrong. The legs and feet are thick and flout, and of nearly the same colour with the bill. The tail is most beautifully barred with deep fearlet and black; the first being the ground or predominant color, on which the latter is disposed in numerous fasciæ. Near the shafts of the feathers on each fide, the fearlet gradually declines into yellow: at the base and tip the tail is wholly black.

The whole bird is of so superior a magnificence as justly to be regarded as one of the finest of its tribe. It is subject to some variation, and in some specimens the scarlet on the tail, instead of being barred with black, forms one large transverse band in the middle.

## BOA CONSTRICTOR.

**◆**\$

CHARACTER GENERICUS.

Scuta abdominalia.
Scuta fubcaudalia (abfque crepitaculo).

Lin. Syst. Nat. p. 373. Cl. Amphibia. Ord. Serpentes.

CHARACTER SPECIFICUS.

BOA VARIEGATA, scutis abdominalibus 240, subcaudalibus 60.

Qui vasta et mirabilia Naturæ opera nunquam automtat contemplati sunt, ii sane quicquid varie de illis
scripserint peregrinatores, caute et dubitanter recipere
solent; immo sæpe utpote sabulosum omnino rejiciunt. Inter hæc Naturæ magnalia jure numerandi
sunt serpentes illi ingentes, qui in nonnullis Indiæ,
Africæ, et Americæ regionibus inveniuntur; quique
in tantam magnitudinem adolescunt ut quadrupedia
etiam majora ingurgitare possint; et longitudine adeo
sunt enormi, ut sæpe pedes viginti, viginti-quinque,
vel etiam triginta superent. Horum temporis progressu multo rarior est conspectus, illosque probabile
est regiones olim sane desertas et nunc excultas, populosque frequentes reliquisse, et in tesqua deserta et

remota exulasse. Spectantur tamen aliquando, hortos et loca habitaculis fuis propiora perreptantes.

Felix fane faustumque est humano generi mirandos hosce serpentes veneno prorsus carere; ob molem tantum et robur timendos; quibus sane adeo præpollent ut cervos aliofque majores quadrupedes, constringendo et amplexando enectos, lento haustu absorbeant: et dein usque ad stuporem et torporem satiati latent, detectique facile vel sclopetis, vel aliis modis confici possint.

Cum prægrandes hos ferpentes quafi ad aliorum animalium perniciem natos contemplamur, horrescamus illos et aversemur necesse est; si vero formam tantum et colores spectemus, non mediocrem illis inesse pulchritudinem lubentissime fatebimur. Color illis, uti et aliis hujus generis, diversus est pro varia ætate; ut plurimum tamen est cinereus, per dorsum et latera plagis maculifque oblongis, magnis, fusco-rubentibus, feu ferrugineis, margine nigro circumfcriptis, et albo flavoque notatis variatus: venter plerumque flavescens albet.

Referunt historici totum Romanum exercitum, conspecto immodico serpente, (quem verisimile est ejusdem speciei fuisse cum illo qui in tabula depingitur), ingenti mețu perculfum fubito, et consternatum suisse. Immanem hunc ferpentem Valerius Maximus ex libro Livii historiæ deperdito his verbis memorat.

" Quæ quia supra usitatam rationem excedentia attigimus, serpentis quoque a T. Livio curiose pariter ac facunde relata fiat mentio: Is enim ait in Africa apud Bagradam flumen tantæ magnitudinis anguem


fuisse ut Attilii Reguli exercitum usu amnis prohiberet; multisque militibus ingenti ore correptis, compluribus cauda voluminibus clisis, cum telorum jactu persorari nequiret, ad ultimum balistarum tormentis undique petitam, filicum crebris et ponderosis verberibus procubuisse; omnibusque et cohortibus legionibus ipsa Carthagine visam terribiliorem: atque etiam cruore suo gurgitibus imbutis, corporisque jacentis pestifero assatu vicina regione polluta, Romana inde summovisse castra: dicit belluæ etiam corium exx pedes longum in urbem missum." Val. Max.


Livium vero ipsum elegantibus supplementis æmulatus est Joannes Frienshemius; audiant igitur velim lectores Frienshemium miram rem suo more diserte

recitanteni.

"Interea M. Regulus proxima quæque subigendo in loca pervenerat per quæ flumen Bagrada labitur: ad quod castra habentes Romanos improvisa pestis et damno non mediocri, et terrore adhuc majore perculit. Magnitudinis enim portentosæ serpens aquatum profectos milites invasit; territisque et nequidquam repugnantibus multos ingentis oris hiatu forbuit: alios spirarum voluminibus et caudæ verbere obtrivit : nonnullos ipfo pestilentis halitus adslatu exanimavit : tantumque negotii M. Regulo facessere potuit, ut totis viribus cum eo de possessione amnis sucrit dimican-Quod ubi cum jactura militum fiebat, neque vinci aut vulnerari dráco potuerat, durissima squamarum lorica, quidquid telorum ingereretur, facile repellente; confugiendum ad machinas, advectifque ballistis et catapultis velut arx quædam munita, dejiciendus

ciendus hostis fuit. Post aliquot jactus in vanum emissos ingens faxum spina dorsi perfracta, vigorem impetumque formidalis monstri resolvit. Sic auoque difficulter confectum est, tanto cum horrore legionum et cohortium, ut se oppugnare Carthaginem, quam alteram talem bestiam malle faterentur. Neque diutius ibi morari castra potuerunt, quin tabo infectas aquas, et omnem circa regionem fætore jacentis pestifero adflatam fugerent. Non fine rubore aliquo humanæ fuperbiæ. quæ non raro viribus fuis nihil negatum effe stolide Exercitum certe Romanum imperatore M. Regulo, terra marique victorem unus anguis et vivus exercuit, et interfectus fubmovit. Quare nec puduit Proconfulem hujus etiam hostis spolia Romam mittere, timorifque fui magnitudinem et victoriæ gaudium publico monumento fateri. Corium enim belluz detractum in urbem devehendum curavit: quod centum viginti pedes longum fuisse dicitur; et in templo quodam suspensum, ad Numantini usque belli tem. pora duravisse." Joan. Friensh. Suppl. Livian.





Landon . - Retiched Book roly by 15:12 land for Soft a Bours level

#### THE

## GREAT BOA.

**◇**◆◆◆<del></del>◆◆<del></del>◆◆<del></del>

GENERIC CHARACTER.

Transverse Lamellæ both on the abdomen and beneath the tail.

SPECIFIC CHARACTER.

VARIEGATED BOA with about 240 Lamellæ on the abdomen and 60 beneath the tail.

By those who are unacquainted with the wonders of Nature, the descriptions given by Naturalists of some of the more striking and singular animals, are received with a degree of cautious scepticism, or even rejected as exceeding the bounds of credibility. Amongst these animals may well be numbered the prodigious ferpents which are sometimes found in India, Africa, and America: ferpents of so great a fize as to be able to ingorge even fome of the larger quadrupeds, and of so enormous a length as to measure 20, 25, and 30 feet. reason to believe that these immense serpents are become less common now than some centuries backwards, and that in proportion as cultivation and population have increased, the larger species of noxious animals have been expelled from the haunts of mankind, and driven into the more distant and uncultivated tracts. They are still however occasionally seen, and sometimes

times approach the plantations and gardens of the diftricts nearest to their residence.

It is happy for mankind that these gigantic serpents are not poisonous; they are therefore to be dreaded only on account of their vast size and strength, which latter is so great as to enable them to kill cattle, deer, and other animals by writhing themselves round them, so as to crush them to death by mere pressure; after which they swallow them in a very gradual manner, and when thus gorged with their prey, grow almost torpid with repletion; and if discovered in this state may without much dissiculty be dispatched by shooting or by other methods.

Confidered as one of the great destroyers of the animal world, this serpent must be viewed with horror; but if we take into confideration only its form and colors, we cannot deny it a very great share of external beauty; like many of its tribe, it is apt to vary in color, and appears in a more or less vivid state according to the period of life in which it happens to be found; but it is generally of a greyish color, variegated on the back and sides in the most curious manner with large oblong patches and spots of bright reddish brown, which are still further decorated by having black margins, surrounded both internally and externally with streaks of white and yellow: the belly is commonly of a yellowish white.

It was in all probability an enormous specimen of this very serpent that once diffused so violent a terror amongst the most valiant of mankind, and threw a whole Roman army into dismay. Historians relate

this surprizing event in terms of considerable luxuriance. Valerius Maximus thus mentions it from Livy, in one of the lost books of whose history it was related more at large.

"And fince we are on the subject of uncommon phænomena, we may here mention the ferpent, fo eloquently and accurately recorded by Livy; who fays that near the river Bagrada in Africa a fnake was feen of so enormous a magnitude as to prevent the army of Attilius Regulus from the use of the river; and after fnatching up feveral foldiers with its enormous mouth, and devouring them, and killing feveral more by flriking and fqueezing them with the spires of its tail, was at length destroyed by affailing it with all the force of military engines and showers of stones; after it had withflood the attack of their spears and darts: that it was regarded by the whole army as a more formidable enemy than even Carthage itself; and that the whole adjacent region being tainted with the peftilential effluvia proceeding from its remains, and the waters with its blood, the Roman army was obliged to remove its station; he also adds that the skin of the monster, measuring 120 feet in length, was fent to Rome as a trophy." VAL. MAX.

The learned Frienshemius, in his Supplementa Liviana, has attempted a more ample and circumstantial narrative of the same event, and it cannot be unsatisfactory to the reader to receive a quotation from an author, who has so happily imitated the manner of the great historian.

"In the mean time Regulus, every where victorious, led.

led his army into a region watered by the river Bagrada, near which an unlooked for misfortune awaited them, and at once affected the Roman camp with confiderable lofs, and with apprehensions still more terrible; for a serpent of prodigious fize attacked the soldiers who were fent for water, and while they were overwhelmed with terror, and unequal to the conflict. engulphed feveral of them in its enormous mouth, and killed others by writhing round them with its spires, and bruifing them with the strokes of its tail: and some were even destroyed by the pestilential essluvia proceeding from its breath; it caused so much trouble to Regulus that he found it necessary to contest the possesfion of the river with it by employing the whole force of his army; during which a confiderable number of foldiers were loft, while the ferpent could neither be vanquished nor wounded; the strong armour of its scales easily repelling the force of all the weapons that were directed against it; upon which recourse was had to battering engines; with which the animal was attacked in the manner of a fortified tower, and was thus at length overpowered. Several discharges were made against it without success, till its back being broken by an immense stone, the formidable monster began to lose its powers, and was yet with difficulty destroyed; after having diffused such a horror amongst the army, that they confessed they would rather attack Carthage itself than fuch another monster. Nor could the camp continue any longer in the fame station, but, was obliged to fly; the water and the whole adjacent region being tainted with the pestiferous essluvia. A most.

most mortifying humiliation to human pride! which fometimes vainly imagines nothing capable of refifting Here at least was an instance of a the powers of man. whole Roman army under the command of Regulus, and univerfally victorious both by fea and land, oppofed by a fingle fnake, which conflicted with it when living, and even when dead obliged it to depart. proconful therefore thought it no diminution to his dignity to fend the spoils of such an enemy to Rome, and to confess at once the greatness of his victory and his terror by this public memorial; for he caused the skin of the fnake to be taken off, and fent to the city; which is faid to have measured 120 feet: it was suspended in a temple, and remained till the time of the Numantine FRIENSH. SUPPL. LIV. war."

## VOLUTA ARAUSIACA.

**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*** 

CHARACTER GENERICUS.

Animal Limax.

Testa unilocularis, spiralis.

Apertura ecaudata, subeffusa.

Columella plicata: Labio Umbilicove nullo.

Lin. Syft. Nat. p. 1186.

CHARACTER SPECIFICUS, &c.

VOLUTA testa alba fasciis transversis fulvis, columella sex-plicata.

VEXILLUM ARAUSIACUM.

Rara hæc et perelegans concha generis Linnæant Volutæ nominati est species. Hanc ipsam ille nullibi descripsit. Communiter illam appellaverunt physici Vexillum Arausiacum. Indica est concha, et præcipue in insula Amboyna reperitur.

Fig. 1. Testa adulta. Fig. 2. Testa junior.

#### THE

## ORANGE FLAG,

OR

## ORANGE-STRIPED VOLUTE.

**◇**◇◇◇◇◇◇◇◇◇◇◇◇◇◇◇◇◇

GENERIC CHARACTER.

Animal refembling a Limax or Slug. Shell unilocular, spiral. Pillar or Column twisted or plaited.

SPECIFIC CHARACTER, &c.

WHITE VOLUTE, with orange-coloured transverse bands, and (generally) six plaits on the column.


LE PAVILLON D'ORANGE.


Argenv. Append.

Knorr. part 5, t. 1. f. 1.

The elegant and rare shell here figured, belongs to the Linnæan genus Voluta, but is not amongst the species described by Linnæus. Its general name amongst collectors is the Orange-Flag. It is an East-Indian shell, and is principally sound in the island of Amboyna.

Fig. 1. The adult shell. Fig. 2. A somewhat younger shell.





### PSITTACUS GLORIOSUS.

<del></del>

#### CHARACTER GENERICUS.

Rostrum aduncum: mandibula superiore mobili, cera instructa.

Nares in rostri basi.

Lingua carnofa, obtufa, integra.

Pedes scansorii.

Lin. Syst. Nat. p. 139.

### CHARACTER SPECIFICUS.

PSITTACUS macrourus fanguineus, dorso nigro maculato, alis caudaque cyaneis.

Psittacus hic formosus plerasque sui generis antecedit pulchritudine et colorum splendore. Ad species macrouras seu longicaudas pertinet. Color ejus generalis est vivide sanguineus, corporis parte inseriore paulum pallidiore. Plumæ dorsi nigræ sunt, rubro cincæ: tectrices alarum minores lucidissime cæruleæ: versus mediam alarum partem color cæruleus magis cyaneus est, non sine virore aliquo obscuriore: super axillas color plumarum niger in maculam transversam nigram apicibus acuminatis discurrit: cauda prælonga ad imum est cuneatior.

Novam Hollandiam incolit hæc avis, et magnitu dine Pfittacum Alexandri æquat, vel paulo fuperat.

Feminæ multo minus vividus est color; dorsumque olivaceo tinctum.

# SPLENDID PARROT.

<del></del>

GENERIC CHARACTER.

Bill hooked. Upper mandible moveable.

Nostrils round, placed in the base of the bill.

Tongue sleshy, broad, blunt at the end.

Legs short. Toes formed for climbing, viz. two forward and two backward.

SPECIFIC CHARACTER, &c.

LONG-TAILED SANGUINE-RED PAR-ROT, with the back fpotted with black, the wings and tail bright-blue.

PENNANTIAN PARROT.

Latham's Synops. Suppl. p. 61.

This beautiful bird exceeds most of its genus in the fplendor of its colors. It ranks amongst the Psittaci Macrouri, or Long-tailed Parrots. Its general color is a vivid fanguine-red, somewhat paler on the lower part of the body; and the feathers on the back are black, margined with red. The shoulder-feathers, or smaller coverts, are of a most brilliant blue: towards the middle part of the wing the blue becomes much deeper, and slightly tinged with green. Above the axillæ the black color of the back forms on each side a transverse





transverse spot with sharpened extremities. The tail is very long, and cunciform towards the tip.

This species is a native of New Holland, and is about the fize of the common Ring-Parrakeet, or rather larger. The semale is much less brilliant in color, and the back is of an olive green.



# STAPELIA HIRSUTA.

**◆\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*** 

CHARACTER GENERICUS.

Corolla contorta.

Nectarium stellula duplex in medio corollæ.

Pentandria Digynia.

CHARACTER SPECIFICUS, &c.

STAPELIA denticulis ramorum erectis.

Lin. Spec. Plantar. p. 316.

ASCLEPIAS AFRICANA AIZOIDES, flore pulchre fimbriato.

Comm. rar. 19. t. 19.

Inter pretiofissimas plantarum dotes merito numeratur odor ille gratissimus quem plurimæ emittunt, quo in colligendo, conservando, augendo, subtiliter studioseque semper laboratum est. Suave-olentium vegetabilium tanta sane est diversitas, ut odores speciatim recensere vix possibile sit. Ut taceam rosam, violam, dianthum, cæteramque copiam narium in hortis plerisque repertam, genera Diosmæ, Illicii, Nyctanthes, et Polyanthes, velut exempla suavissimæ et pollentissimæ fragrantiæ satis sit proferre.


Huic aromaticorum ubertati opponuntur exempla aliquot contrarii odoris in generibus Ari, Sterculiæ, Lobeliæ, cæterifque nonnullis; rara hæc quidem, quæ tamen fatis probant Naturam in humani generis oblectatione

lectatione non femper versari. Phalli species, in sylvis, exeunte æstate, non raro conspecta, odorem quasi carnis putridæ spargendo, insecta multa carnivora, muscas præcipue vulgares, seu carnarias ad se allicit, quæ in summitate plantæ catervatim conglomerantur, liquorem glutinosum exinde copiose essuentes.

Omnes autem hactenus cognitas, quæ more mortui cadaveris fætere folent, planta illa in tabula depicta longe superat; cui tanta est odoris putridi graveolentia, ut e longinquo veniant muscæ vulgares seu carnariæ Linnæi, ovaque ibi deponant; quibus exclusis, larvæ parvo tempore plerumque pereunt; qua in re unica feipfam decipere, fuaque quafi frustrare confilia videatur Natura. Interdum tamen, si qua sides observatoribus, accidit ut aliæ muscarum species, carnaria multo minores, in eodem flore ova fua deponant, larvæque excluíæ, ob exiguitatem parvo cibo egentes, florem ipfum depafcant; ad plenam magnitudinem perveniant, et in chryfalides convertantur, ex quibus muscæ persectæ postea erumpunt. Ita se rem habere demonstrat experimentum a domino Hill memoratum, qui narrat larvas plurimas muscarum, hoc solo flore enutritas, omnibus vicibus et permutationibus absolutis, ad plenam et perfectam maturitatem adolevisse.

Africæ est indigena hæc planta, et in promontorio bonæ spei præcipue invenitur.





# ·FETID STAPELIA,

### OR THE

### CARRION-FLOWER.

**◇**\*\*\*\*\*\*\*\*\*\*\*\*\*

#### GENERIC CHARACTER.

Corolla slightly twisted towards one side.

NeEtary in form of a double star in the middle of the corolla.

Pentandria Digynia.

SPECIFIC CHARACTER, &c.

STAPELIA with the denticles of the branches upright.

THE CARRION-FLOWER.

The fragrance emitted by the major part of the vegetable world is such as to be regarded amongst the most elegant and valuable attributes of plants; and the ingenuity of mankind has been studiously employed in collecting, preserving, and concentrating it. So great is the variety of vegetable persumes that it is hardly possible to enumerate the different kinds. Without particularizing the rose, the violet, the carnation, and many other common inhabitants of the garden, we might adduce the genera of Diosima, Illicium, Nyctanthes, Polyanthes, and some others, as peculiar

peculiar examples of the richest and most exalted fragrance.

Amidst this profusion of sweets in the vegetable kingdom there are occasionally interspersed some examples of a contrary odor: thus some species in the genera of Arum, Sterculia, Lobelia, and a few others, are convincing proofs that Nature does not always accommodate her productions to the immediate gratisfication of the human sense.

A fpecies of Morell, not uncommon in woods towards the latter part of the fummer, has a fmell fo much refembling that of putrid animal matter, as to attract feveral species of carnivorous infects, and particularly flies, which are generally observed fettled in great numbers on the top of the plant, and eagerly feasting on the glutinous moisture with which it abounds.

of a finell fimilar to that of animal fubflances, the plant here figured is by far the most remarkable; and when in a state of florescence, disfuses a scent so powerfully cadaverous as to allure the common sless-fly, or blow-fly to deposit its eggs on the flower, where they are hatched, and the young larvæ, for want of proper nutriment, commonly perish soon afterwards; and thus Nature in this single instance, almost seems to have deceived herself. If however we may rely on some observations, it happens sometimes that other species of slies much smaller than the common sless-fly, deposit their eggs in the slower, and when hatched, the larvæ being very small, and not requiring much sood,

do actually find a fufficient nutriment from the flower itself, till the time of their change to chrysalis arrives; and after the usual period become complete flies; and this appears to have been the case in an instance recorded by the late Sir John Hill, where several larvæ of flies are said to have passed through all their changes, though nourished solely by the flower of this plant.

It is a native of Africa, and is principally found at the Cape of Good Hope.



# SOREX BICOLOR.

�\*\*\*\*\*\*\*\*\*\*\*\*

CHARACTER GENERICUS.

Dentes Primores superiores duo bisidi.

Inferiores quatuor: intermediis breviori-

bus.

Laniarii utrinque plures.

Lin. Syft. Nat. p. 73.

CHARACTER SPECIFICUS, &c.

SOREX cauda mediocri, corpore fupra nigricante, fubtus albicante.

SOREX cauda mediocri fubnuda, corpore nigricante, fubtus cinereo.


Erx. Mamm. p. 124.

MUSARANEUS dorso nigro ventreque albo.

Merret. pin. p. 167.

Inter quadrupedes Britanniæ rariores merito numeratur parvulus hic Sorex, ab illis folummodo qui præcipuam scientiæ naturali navant operam, observari solitus. In Europa sere omni nascitur, et ut docet dominus Pallas, in Sibiria frequenter reperitur. In Gallia annum circiter millesimum septingentesimum quinquagesimum sextum a domino Daubenton detectus et descriptus est. In Anglia longe antea cogni-

tus, inter animalia tamen deperdita habebatur, donec annum circiter millesimum septingentesimum sexagesimum octavum, juxta cœnobium de Reevesby in comitatu Lincolniensi iterum deprensus sit. Vix dubitandum est quin in aliis multis Angliæ partibus possit in-Cur tamdiu incognitus manserit in causa esse videtur obscuritas latibulorum, (amat enim ripas fluviorum, locaque aquofa,) nec non communis hujus-In certis Buckingamiæ locis modi rerum incuria. nequaquam rarus est, et Oxoniæ vicinio: sedem vero fibi potissimum delegit in aggeribus fluminis Collegii Beatæ Mariæ Magdalenæ ambulacra circumeuntis; ubi non raro mortuus jacet, quod et Sorici Araneo LINNÆI, seu vulgari sæpius accidit, a qua specie sacillime primo intuitu, colore nempe longe diverso, dignosci potest. 4



#### THE

### WATER-SHREW.

**\*** 

GENERIC CHARACTER.

Two cutting Teeth in each jaw. Long flender Nofe: fmall Ears. Five Toes on each foot.

Pennant.

SPECIFIC CHARACTER, &c.

SOREX black above, whitish beneath.

MUSARAIGNE D'EAU.

Daubenton Mem. de l'Acad. de Paris, 1756, p. 211. t. 5. f. 2.

WATER-SHREW.

Pennant Hist. Quadr. p. 480.

This animal may be numbered amongst the rarer quadrupeds of Britain, and is but very seldom noticed except by those who pay a more than ordinary degree of attention to the natural productions of our kingdom. It appears to be a native of almost every part of Europe, and from the observations of Dr. Pallas it should seem to be very well known in Siberia. In France it was discovered and described by Mr. Daubenton in the year 1756. In England it had long before been noticed;

ticed: but was confidered amongst the lost quadrupeds of our island till about the year 1768, when it was again observed in the neighbourhood of Reevesby-Abbey in Lincolnshire. It may probably be discovered in many other parts of the kingdom, and its continuing fo long undiffinguished must partly be ascribed to the obscurity of its retreats, under the banks of rivers, and in watery places, and partly to a mere want of attention to objects of this nature. In some parts of Buckinghamshire it is not very uncommon, it is also found in the neighbourhood of Oxford; and it feldom fails to refide in the banks of the river furrounding the water-walks of Magdalen College; and is not unfrequently found dead by the edges of the walks, in the fame manner as the common Shrew, or Sorex Araneus of Linnæus, from which it may always be most readily distinguished by its very different color.

• • 1



# LOXIA JUGULARIS.

<u></u>

#### CHARACTER GENERICUS.

Rostrum conico-gibbum, frontis basi rotundatum versus caput: Mandibula inferior margine laterali inflexa.

Nares in basi rostri.

Lingua integra:

Lin. Syst. Nat. p. 299.

#### CHARACTER SPECIFICUS.

LOXIA testaceo-ferruginea, nigro undulata, fascia gulæ fanguinea.

Africam incolit hæc avicula. Coloris est Castanei; seu griseo-serruginei, maculis lunatis nigris eleganter undulati: Decoratur gula fascia lata transversa vivide coccinea.

#### THE

# RED-THROATED GROSSBEAK.

**◇**◇<u>♦</u>◆<u>\$</u>◆<del>\$</del>◆<del>\$</del>

GENERIC CHARACTER.

Bill strong, thick, and convex.

Nostrils small.

Tongue truncated.

SPECIFIC CHARACTER, &c.

PALE-FERRUGINOUS LOXIA, undulated with black, with a blood-red fascia on the throat.

FASCIATED GROSSBEAK.

Lath. Synopf. 2. p. 156

THE CUT-THROAT SPARROW.

Lev. Muf.

This bird is a native of Africa. Its colour is a palish chesnut, or grey ferruginous, elegantly undulated with semi-lunar marks of black. The throat is ornamented with a broad transverse band of the deepest vivid red.





## CHÆTODON ARMATUS.

**•** 

### CHARACTER GENERICUS.

Dentes fetacei, flexiles, confertissimi, numerosissimi.

Membrana branchiostega radiis sex.

Corpus plerumque fasciatum. Pinna dorsi anique carnosa squamosa.

### CHARACTER SPECIFICUS.

CHÆTODON ALBESCENS, corpore fasciis septem nigris, spinis pinnæ dorsalis sex, tertia longissima.

Novam et perelegantem generis Chætodontis speciem dat tabula inspiciendam. Vix quatuor uncias superabat speciminis longitudo unde hæc nostra sigura delineata est. Color huic Chætodonti est argenteo-albus, dorso sub-cæruleo. Fasciæ corporis transversæ nigerrimæ sunt: pinnæ caudaque pallide susca: spina seu radius tertius pinnæ dorsalis anterioris cæteris multo longior. Maris Australis incola est hic piscis; prope novam Hollandiam captus est, indeque nuperis navigiis ad nos allatus.

#### THE

# LONG-SPINED CHÆTODON,

### · GENERIC CHARACTER.

Teeth fetaceous, flexile, closely fet, extremely numerous.


Branchiostegous membrane with fix rays.

Body generally fasciated. Dorsal and anal Fins fomewhat sleshy and coated with scales.

#### SPECIFIC CHARACTER.


WHITISH CHÆTODON, with feven black bands on the body, fix spines on the dorsal fin, the third very long.

This fish is a new and very elegant species of the genus Chætodon. The total length of the specimen from which the figure was taken was scarcely more than four inches: the color is a silvery white; darker, and of a blueish tinge on the back: the transverse safciæ or bands are of a deep black: the fins and tail are of a pale brown: the third ray or spine of the first dorsal sin is much longer than the rest. It is a pative of the Southern Ocean, and was brought from New-Holland in one of the last voyages.



Landon . Bellistad istant of for the for by Sind at the Con at 12 a Brown Source

J. .



# BRADYPUS URSINUS.

CHARACTER GENERICUS.

Dentes Primores nulli utrinque. Laniarii obtusi, folitarii, molaribus longiores. Molares utrinque 5 vel 6.

CHARACTER SPECIFICUS.

BRADYPUS niger hirfutiffimus, nafo elongato nudo.

Inter plurima quæ phyficis nuperrime innotuerunt, principem fibi locum non immerito vindicat animal in tabula depictum; quod fortaffe nonnullis videatur nowum omnino per fe genus conflituere. Opinione tamen firmatus celeberrimi Pennanti, cujus comes, in illius characteres primum fedulo inquifivi, ad genus Bradypi referendum cenfeo.

Habitu corporis externo ad ursum vulgarem tam prope accedit Bradypus Ursinus, ut pro Ursi specie a quibusdam habitus sit. Vetant tamen omnino dentes, unguesque peculiariter sormati, cum cæteris characteribus, ut Ursi speciem esse pronunciom.

Magnitudine est ursi vulgaris, exceptoque vultu, seu rostro, villo nigro, longo, et hirsuto totus obtegitur; qui tamen super collum et tergum multo longior est quam in cætero corpore. In partibus corporis anteriporibus antrorsum spectat hirsutia; in posterioribus retrorsum.


trorfum. Oculi minimi: aures parvæ, et denfo capiris vellere fere absconditæ. Dentes incisores nulli sunt Utrique maxillæ insunt duo dentes laniarii modici: qui in maxilla superiore siti sunt uncias circiter duas a se invicem distant; qui in inferiore, paulo miuus. maxilla superiore insunt dentes molares tres utrinque, quorum duo maxime remoti feu interiores funt quasi duplices et lobati; alter fimplex est. In inferiore funt utrinque sex molares, quorum remotissimus seu interior fimplex est; duo huic proximi duplices, cæterique tres fimplices. Linguæ lævi nihil peculiare est. Rostrum elongatum est, et tensile, videturque cartilagine transversa et mobili interius esse instructum. Ungues pedum anticorum, qui quinque, sunt validissimi, modice incurvati et acuminati: pedum posteriorum ungues breviores funt et potius rotundati : Cauda brevissima vix distingui potest.

Ab interiore Bengala allatus est Bradypus Ursinus, et quartum jam annum excessit. Primo detectus Ursum Lotorem Linnæi, qui Raccoon vulgo dicitur, magnitudine vix superavit; illumque latratum quasi caninum emissise asserunt custodes. Vox illi nunc nulla est præter rugitum brevem et abruptum, qui non nisi lacessito et irritato prodit. Mitis est et tranquilli ingenii. Vegetabilibus et lacticiniis vescitur: gaudet pomis, carnesque fastidit, nisi forte tenerrimas. Medullam tamen ex osse lubentissime exugit. Melle, saccharo, et omnigenis cupediis delectatur. Motus huic non ut in aliis ejusdem generis, tardissimus, sed modicus. Solet itetum atque iterum, more canis cubitum ituri, circumgyrare.


gyrare. Terram scalpere, et sub humo velle se occultare dicitur, unde a repertoribus primo essossus est.

Ab ingeniofo Domino Catton figuram hanc noftram mutuati fumus, qui quicquid animali proprium et fingulare est optime expressit.

In altera tabula ringens exhibetur caput, ut fitus dentium oftendatur, nec non ut roftrum fuperius, ope cartilaginis internæ egregie flexibile, quodammodo demonstretur.











# ÚRSINE BRADYPUS,

#### URSIFORM SLOTH.

GENERIC CHARACTER.

Cutting-Teeth none in either jaw.

Canine-Teeth folitary, longer than the grinders.

Grinders in each jaw 5 or 6.

SPECIFIC CHARACTER.

BLACK BRADYPUS with very long fhaggy hair, and naked lengthened fnout.

Amongst the new species of animals with which Zoology has been lately enriched, the creature here sigured claims a distinguished place; and it may be doubted whether it might not with great propriety constitute a new genus. In compliance however with the opinion of Mr. Pennant, in whose company I first made an accurate examination of its characters, I have here placed it as a species of the genus Bradypus, or Sloth.

In its habit, or general appearance, it has a striking resemblance to the common bear, and it has even been considered as a species of bear by some naturalists.

C c

Its teeth, however, and the peculiar formation of its claws, with feveral other particulars, absolutely forbid it to be any longer confidered as a species of Ursus.

The animal is about the fize of a bear, and is covered all over, except on the face, or rather the fnout, with long, fhaggy, black hair, which on the neck and back is much longer than in other parts. On the fore part of the body the hair points forwards, and on the hinder part backwards. The eyes are very fmall; the ears rather small, and partly hid in the long hair of the head. It is totally destitute of incisores or frontteeth: in each jaw are two canine teeth of a moderate fize: those in the upper jaw are situated at the distance of two inches from each other: those in the lower jaw are placed fomewhat less remote. The molares or grinders in the upper jaw are three on each fide; of which the two most remote are double, or lobed: the remaining one fimple. In the lower jaw there are on each fide fix grinders, of which the most remote or backward is simple; the two next double, and the three remaining ones fimple. The tongue is fmooth, and has nothing remarkable in its appearance. The nofe, or fnout is of a somewhat elongated form; it also appears as if furnished with a fort of transverse joint, or internal cartilage, which admits of a peculiar kind of motion in this part. The claws on the fore-feet are five in number, and are excessively strong, moderately crooked, and fharp pointed: those on the hind-sect are shorter, and of a rounder shape. The tail is very thort, and inconspicuous.

This animal was brought from the interior parts of Bengal, and is now fomewhat more than four years old. When first taken, it is said to have been about the size of a raccoon, and is reported to have fometimes barked in the manner of a dog. Its voice however at prefent is rather a fort of fhort, abrupt roar, which it emits when much disturbed or irritated. It is gentle and good-natured: feeds chiefly on vegetable substances and milk; and is fond of apples, and does not willingly eat animal food, except of a very tender nature, as marrow, which it readily fucks from a bone prefented to it. It is also delighted with honey, sugar, and other fweets. Its motions are not as in others of this genus, flow and languid, but moderately lively, and it appears to have an habit of turning itself round and round every now and then, as if for amusement, in the manner of a dog when lying down to fleep. It is faid to have a propenfity to burrowing under ground; and that it was at first dug out of its retreat by those who discovered it.

The figure here given is copied from a drawing by Mr. Catton, an artist who has been peculiarly happy in expressing the appearance of the animal.

In the additional plate is represented a view of the head in a ringent state; in order to shew not only the form and disposition of the teeth, but also in some measure the singular stexibility of the upper part of the snout, as if surnished with a joint or internal cartilage.



## PETAURUS AUSTRALIS.

CHARACTER GENERICUS.

Dentes varii in variis. Corpus cute volatili cinctum.

CHARACTER SPECIFICUS.

PETAURUS fupra nigricans, fubtus albicans, pollicibus plantarum rotundatis.

Arbitrentur fortasse Physici debere Petaurum Australem genus potius per se constituere, quam inter novas species numerari. Cum vero generum institutio ob hanc vel illam peculiarem notam, pedum nempe seu dentium formam, scientiam naturalem sæpe perturbet et obscuret, molestiamque potius quam auxilium tyroni afferat ; fatius duxi Petaurum Australem velut speciem Sciuri volantis describere. Sciurosque d lpsos volantes ab aliis disjungere, et in genere distincto nomine Petauri reponere. Quamvis enim a Linnæo in eodem genere cum Sciuris aliis numerentur, fatendum tamen est, membranam, seu cutim extensam, cujus ope volitant, adeo esse singularem et propriam, ut non immerito ad genus separatum possint pertinere.

Hoc tamen fortasse nonnullis videatur curiose potius quam utiliter factum.

Totius igitur generis Petauri, species hic depicta non tantum maxima est, sed et elegantissima. præterca Dd

præterca est nova, de Nova Hollandia non ita pridem petita. Magnitudine cuniculo juniori vix cedit.

Color corporis fuperni est susco-cinereo-nigricans, inferioris fere albidus. Pilus est præter modum mollis et elegans; cujus si copia sufficeret, nescio annon aliorum animalium vellera, utcunque pretiosa, sordescerent. Pedum posticorum forma singularis est: Pollex seu digitus primus rotundatus est, unque complanato et rotundato; cum aliis omnibus digitis unques acuminati sint. Digiti quoque duo intermedii, seu pollici proximi, cute communi ad ipsos unques conjunguntur. Cauda longa pilo mollissimo et delicatulo densissime vestitur.

In narratione itineris australis a Domino White, hoc animal nomine *Hepoona Roo* distinguitur, et pulcherrime a Domino Catton depingitur: figuram igitur ibi datam, utpote eximiam, in hoc nostro opere imitamur.





buston , Robbished March stongs by Till Vollor of the A. to Bourse Stoner.

#### THE

## SOUTHERN PETAURUS.

Ø\$\$\$\$\$\$\$\$\$\$

GENERIC CHARACTER.

Teeth differing in the different species. To be Body surrounded by a flying-skin.

SPECIFIC CHARACTER, &c.

PETAURUS blackish above, whitish below, with the thumbs of the hind-feet rounded. HEPOONA ROO.

White's Voyage to Botany Bay.

It may perhaps be doubted whether this animal might not conflitute a genus per se, rather than be regarded as a species merely: but as the institution of genera from two or three particular circumstances only; e.g. a peculiarity in the teeth, claws, &c. frequently throws much confusion on natural history, and rather tends to embarrass than instruct a student in the science, I have thought it best to describe this animal as a species of slying-squirrel, and to separate the slying-squirrels from the genus Sciurus, with which Linnæus had conjoined them, and to form them into a genus by the name of Petaurus.

This perhaps may be thought an unnecessary piece of exactness; yet there is something so peculiar in the expanded processes of skin by which the slying-squir-

Dd 2

rels are distinguished, that they may properly enough

Of all the species then of this genus the animal here figured is the largest and the most elegant: it is also a new species; having been brought from New Holland during the late voyages.

Its color-as a very fine fable or deep grey-brown above, while the lower furface is nearly white. The fur is of the most exquisite softness and elegance, and if the animal could be procured in sufficient plenty, would bid fair to supersede the richest surs hitherto in use. The structure of the hind foot in this species is remarkable; the thumb or first toe being of a rounded form and surnished with a flattened nail, while the remaining toes have sharp claws; and the two second toes, or those next the thumb, are united in one common covering or skin to the very claws: the tail is long, and is very thickly cloathed with sur of the softest and most delicate appearance.

In Mr. White's Journal of a voyage to New South Wales, this animal is diffinguished by the title of Hepoona Roo, and is beautifully figured by Mr. Catton; which figure, on account of its excellency, we have not scrupled to copy in the present work.

## SIREN LACERTINA.

CHARACTER GENERICUS.

Corpus bipedum, caudatum, nudum. Pedes brachiati, unguiculati.

Lin. Syst. Nat. vol. 1. pt. 2. Addend.

CHARACTER SPECIFICUS, &c.

SIREN corpore anguilliformi, branchiis ramofis. SIREN LACERTINA.

Amæn. Acad. vol. 7. p. 311.

Sirenem a reliquis animalibus adéo eminenter distinxit Natura, forma ambigua, et dubiis characteribus, ut celeberrimus Linnæus novum illi Amphibiorum ordinem nomine *Meantes* instituerit; qui tamen novus ordo non inter alia Amphibia in Systemate Naturæ invenitur, sed in nota ad finem partis secundæ tomi primi istius operis exstat.


Genus cui Siren affinitate summa conjungi videtur, est genus Lacerta: lacertæ enim larvam valde repræfentat, et dubitant nonnulli an talis revera sit nec ne: characteres tamen omnes videntur esse animalis perfecti et adulti, nec hactenus visa est Siren in alia aliqua forma statuque. Pedes tantum duos habet, sine ullis pedum posteriorum vestigiis; illosque unguibus munitos, cum larvarum omnium lacertarum, hactenus cognitarum, pedes sunt, (lingua Linnæana) mutici, sive unguibus


unguibus destituti. Os dentium parvulorum seriebus pluribus munitur.

Singulare hoc animal Americam Septentrionalem inhabitat, et in Carolina Australi præcipue invenitur, in locis uliginosis et aquosis minime infreques, sub aqua plerumque degens, interdum vero in terram progrediens. Vocem habet stridulam, sive cantillantem, quam ob causam a Linnæo nomine Sirenis distinguitur.

A Domino Garden, medico in Carolina per multos annos celeberrimo, primo detectum fuit descriptumque; qui etiam specimina nonnulla ad Linnæum mifit. Linnæus in epistola sua ad Dominum Garden de Sirene, affirmat nullum aliud unquam animal se adeo torsisse, nec aliud se plus velle, quam veram istius naturam optime callere. Non silendum est Sirenem in terram modica vi jactatam, in partes tres vel quatuor dissilire, quod aliquibus etiam serpentibus evenit.

In Actorum Anglicorum volumine quinquagesimo sexto describitur Siren a Domino Ellis, qui etiam siguram addidit, quæ adeo ad normam animal exprimit, ut potius visum sit eam ipsam imitari, quam novam, diverso situ, essingere. Summam Sirenis larvæ lacertæ similitudinem demonstrat triplex branchiarum ramosarum utrinque in collo par, ut sit in larva lacertæ vulgaris aquaticæ. Species lacertæ cui Siren maxime videtur assinis, est lacerta a Linnæo Teguixin nominata, Ad longitudinem sere duorum pedum crescit Siren.





Landers, Retracted Mand 1979, by Soil , Soilder Soff & to Bours lines .

#### THE

### SIREN.

<del>\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*</del>

#### GENERIC CHARACTER.

Body two-footed, tailed, naked.

Feet brachiate (i. e. placed in the manner of arms), and furnished with claws.

#### SPECIFIC CHARACTER.

SIREN with an eel-shaped body, and ramified branchiæ, or respiratory organs.

The Siren stands eminently distinguished in the list of animals by the ambiguity of its characters, which are such as to have induced the great Linnæus to institute for it a new order of Amphibia, under the title of Meantes; an order however which does not stand amongst the Amphibia in the Systema Naturæ, but is mentioned in a note at the end of the second part of Vol. I. of that work.

The genus with which the Siren has evidently the greatest possible affinity, is that of Lacerta or Lizard. It even very much resembles the larva, or first state of a lacerta; and it is still doubtful whether it may not really be such; but it has never been observed in any other state, and it has always two seet only, without any appearance of a hind-pair. It likewise has all the characters of an animal in its complete state; the seet in particular, are armed with claws, whereas the larvæ of all the lacertæ are without claws, or in the Linnæan phrase, digitis muticis. The mouth has several rows of smallish teeth.

This most singular animal is a native of North America, and the part where it has principally been found is the province of South Carolina, where it is not uncommon in muddy and swampy places, living generally under water, but sometimes appearing on land. It has a fort of squeaking or singing voice, for which reason Linnæus has applied to it the name of Siren.

It was first discovered by the ingenious Dr. Garden, who resided long in South Carolina, and who sent an account of the animal, accompanied with specimens, to Linnæus. Linnæus in his letter to Dr. Garden on this subject, declares that nothing had ever exercised his thoughts so much, nor was there any thing he so much desired to know as the real nature of this extraordinary creature. It is remarkable that the Siren, when thrown on the ground with a degree of violence, breaks in three or sour pieces; in which particular it resembles some of the servent tribe.

In the fifty-fixth Volume of the Philosophical Tranfactions is an account of the Siren by the late Mr. Ellis, illustrated by a figure which so very accurately expresses the animal, that we have not scrupled to copy that figure, rather than to give a new one in a different posture. What causes this animal to approach very nearly to the appearance of the larva of a lacerta, is, that it is surnished on each side the neck, with three pair of ramissed branchiæ, in the same manner as the larva of the common water-newt. The species of lacerta to which it seems most allied is the Lacerta Teguixin of Linnæus. It grows to the length of nearly two feet.

## GRYLLUS MIGRATORIUS.

**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*** 

#### CHARACTER GENERICUS.

Caput inflexum, maxillosum, palpis instructum. Antennæ setaceæ, seu filisormes.

Alæ quatuor, deflexæ, convolutæ: inferiores plicatæ.

Pedes postici saltatorii. Ungues ubique bini.

Lin. Syst. Nat. p. 693.

### CHARACTER SPECIFICUS, &c.

GRYLLUS thorace subcarinato: segmento unico, capite obtuso, maxillis atris.

Lin. Syft. Nat. p. 700.

GRYLLUS thorace subcarinato: segmento unico, mandibulis cæruleis.

Fabr. Spec. Inf. tom. 1. p. 365.


Præter omnia insecta quæ fruges devorando hominum miserias augere solent, Grylli, sive Locustæ ut communiter dicuntur, maxime metuendam stragem nobis inserunt. In calidioribus terræ partibus, Africæ præcipue et Orientis regionibus, edacissimorum horum animalium innumeræ exurgunt catervæ, quæ supra sidem omnia devastant; tractus per quos volitant, sere desolant, et spatio brevi amænissimas et sertilissimas provincias in deserta convertunt; dum densissimo co-

rum agmine ipse dies adimitur. Dubitari pene possit, fintne locustæ nocentiores vivæ an mortuæ; e mortuarum enim millibus tanta exoritur putredo, ut incipientis pestis una e causis valentissimis censeatur. Locustarum plurimæ funt species, quarum quæ maxime fortaffe vorax et exitiofa est ca in tabula depicta, quæ Gryllus Migratorius LINNÆI. Plerumque coloris est subfusci, sed maculis obscurioribus variati; et in quibusdam corporis partibus coloris est rubentis, seu potius carnei; crura autem plerumque funt subcærulea. Anno millefimo feptingentefimo quadragefimo octavo, in varias Europæ partes irrupit hæc species, sed in catervis sparsis et diffractis. In Germaniam, Galliam, et Angliam involabant, et in urbe ipsa Londini ejusque vicinio plurimæ reperiebantur hujufmodi locustæ: brevi tamen spatio periere, vix ulla agris illata injuria. Maxima omnium specierum hactenus cognitarum est Gryllus cristatus Linnæi, qui insectum jam descriptum mole quincuplici vel etiam fextupla fuperat; quique cum aliis majoribus hujus generis, in nonnullis Orientis regionibus pro cibo usurpari solitus est. Non tantum recentes, sed et sale conditæ comeduntur hæ locustæ, et in foro publico venales prostant in oppidis Levantinis. Substantia tamen earum pergracilis est, præcipue infectorum marium; nam fæminæ ob ovarium paulo pleniorem escam suppeditant. Notissimum est multum disputasse theologos de loco in scriptis evangelicis, ubi dicitur Joannem Baptistam locustis et melle fylvestri vitam sustinuisse, quorum nonnulli per verbum angidas, teneras tantum plantarum summitates intelligi voluerunt. Cum tamen probe compertum

pertum fit, Arabas hisce animalibus vel nunc pro cibo uti, vix egere videtur illa vox ulla alia interpretatione præter communem et vulgarem. Cur enim miremur fanctum illum et abstemium prophetam, cum ab inanibus mundi gaudiis se procul subduceret, cibo usum suisse non fane inter dapes lautiores numerando, sed illo tantum qui in locum potioris escæ suffectus sit? Ut hanc conjecturam firmemus, testimonium Domini Has-SELQUIST liceat nobis proferre, qui de hac ipfa re ita loquitur. " Qui infectis virum hunc fanctum vesci folitum fuisse negant, aiunt cibum e locustis paratum ignotum prorsus esse et inusitatum. Si vero hi in Ægyptum, Arabiam, aut Syriam migrarent, longe aliter de hac re sentirent: certis enim temporibus locustas assatas communiter comedunt Arabes, nec dubitandum est illis tempore quo vixit Joannes Baptista, pro cibo fuiffe. Præterea his locis mores antiqui non facile dilabuntur, nec Baptistæ cibus mirus aut infolitus jam reputatur. Presbyter etiam Græcus, nequaquam indoctus, mihi pro vero assirmavit ecclesiam suam per verbum axeidas locustas semper intellexisse; risitque ipse cum a me audiret nostros interpretes plantam aut avem illo verbo fignificari conjecisse."

entro promotive of main, and the entropy of the ent





#### THE

## MIGRATORY,

OR,

### WANDERING LOCUST.

**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*** 

#### GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with palpi or feelers.

Antennæ either setaceous, or filiform.

Wings four, deflected, convolute; the inferior ones plaited.

Hind-Feet formed for leaping. Claws on all the feet double.

#### SPECIFIC CHARACTER.

GRYLLUS with fubcarinated thorax with a fingle fegment, obtufe head, and blackish or blueish jaws.

Of all the infects which are capable of adding to the calamities of mankind by devouring the products of the earth, locusts seem to possess the most formidable powres of destruction. Legions of these voracious creatures are from time to time produced in various parts of Africa and the Eastern World, where the havock they commit is almost incredible. They carry desolation with

with them wherever they pais, and in the short space of a few hours change the most fertile provinces into the appearance of a barren defart; while the air is darkened by their number's. Nay even when dead they are still terrible, fince the putrefaction which arifes from their inconceivable number, is fuch that it has been regarded as one of the principal causes of a beginning pestilence. There are a great many different fpecies of locusts. One of the most destructive is the Gryllus Migratorius of Linnæus, or the Wandering . Locust, here represented. It is generally of a brownish color, varied with spots of a deeper cast, and in some parts with pale red or flesh-color, and the legs are commonly blueish. In the year 1748 this species appeared in irregular flights in feveral parts of Europe; as in Germany, France, and England, and in London in particular, and its neighbourhood, great numbers were feen: they perished however in a short time, and were not productive of any mischief. The largest species of locust yet known is the Gryllus Cristatus of LINNÆUS, which is five or fix times as large as the species just described; and which, together with some other of the larger kind, is made use of in some parts of the world as an article of food. They are eaten both fresh and salted, in which last state they are publickly fold in the markets in some parts of the Levant. The quantity of edible substance which they afford is but fmall, especially in the male insects; but the females, on account of the ovaries, afford a more nutritions fustenance. It is well known that different interpretations have fometimes been given of the passage

in the facred writings in which John the Baptist is faid to have fed on locusts and wild honey; and the word angidas has been supposed to mean the young shoots of vegetables, rather than locusts: but fince the fact is well established, that these insects are still eaten by the inhabitants of the East, there feems not the least reason for admitting any other interpretation than the usually received one: Why should we wonder that the abstemious prophet, during his state of solitary seclusion from the commerce of the world, should support himself by a repast which is to be numbered not amongst the luxuries of life, but merely regarded as a fubflitute for food of a more agreeable nature? We may also adduce, in support of this idea, the testimony of HASSELQUIST, who thus expresses himself on this very fubject. "They who deny infects to have been the food of this holy man, urge, that this infect is an unaccustomary and unnatural food; but they would foon be convinced of the contrary, if they would travel hither, to Egypt, Arabia, or Syria, and take a meal with the Arabs. Roafted locusts are at this time eaten by the Arabs at the proper feafon, when they can procure them; fo that in all probability this dish has been used in the time of ST. JOHN. Ancient cuftoms are not here subject to many changes, and the victuals of Joнn are not believed unnatural here; and I was affured by a judicious Greek priest, that their church had never taken the word in any other fense, than that of locusts; and he even laughed at the idea of its being a plant or a bird."

Hasselquist's Travels, Eng. Translation, p. 419.



## MUSCICAPA PORPHYROBRON-CHA.

<del></del>

#### CHARACTER GENERICUS.

Rostrum subtrigonum, utrinque emarginatum, apice incurvo; vibrissæ patentes versus sauces. Nares subrotundæ.

Lin. Syst. Nat. p. 324.

CHARACTER SPECIFICUS.

MUSCICAPA nigra, gula purpurea.

Rara hæc avis Americam Australem, et præcipue Cayanam inhabitat. Nigra est, excepta gula, seu colli parte anteriore egregie purpurea. Femina penitus nigricat. Magnitudine merulam vulgarem sere æquat.

#### THE

## PURPLE\_THROATED FLY\_CATCHER.

**◇**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*

#### GENERIC CHARACTER.

Bill fomewhat trigonal, emarginated on each fide, with incurved tip; furnished with spreading vibrisse or whiskers towards the mouth.

Nostrils roundish.

SPECIFIC CHARACTER, &c.

BLACK FLY-CATCHER, with purple throat.


PURPLE-THROATED FLY-CATCHER.

Lath. Synopf. 4. p. 365.

PIAUHAU.

Buff. Oif. 4. p. 588.

This rare and curious bird is a native of South America, and is principally found in Cayenne. It is entirely black, except on the fore-part of the neck, which is of a rich and deep purple. The female is totally black: its fize is nearly that of a black-bird.



5.1.

63



### ACARUS VEGETANS.

#### CHARACTER GENERICUS.

Pedes octo.

Oculi duo ad latera capitis.

Tentacula duo, articulata, pediformia.

Lin. Syst. Nat. p. 1022.

#### CHARACTER SPECIFICUS.

ACARUS crustaceus rufus marginatus convexus, subtus planus.

- Lin. Syst. Nat. tom. 3. p. 226.

Quod physicis haud omnino ingratum esset me facturum arbitratus sum si insectum illud singulare a Linnæo Acarus Vegetans nuncupatum, ob exiguitatem antehac pene neglectum accurate describi et depingi curarem. Aliorum insectorum corporibus plerumque se assigit hujusmodi acarus, illorum præcipue quæ Coleoptera Linnæi constituunt, quæque communiter Scarabæi dicuntur.

Sæpissime illum comperi corpori aut elytris adhærentem Histeri Unicoloris Linnæi, parvi nempe insecti coleoptrati, in hortis, ineunte vere, minime rari. Frequenter etiam eum vidi scarabæo stercorario Linnæi adjunctum, et altero Scarabæo minori, susce nempe varietati simetarii ejusdem auctoris, mensibus Aprilis et Maii, in simo bovino ut plurimum obvii.

Mirandus

Mirandus omnino est modus quo adhærent hi acari corporibus aut elytris aliorum animalculorum, pedicello nempe, seu stipite slexili; quo sit, nisi succus nutritius ab hoc pedicello absorbeatur, (quod ut siat tubulatum putemus,) necesse esse stipitem aliquando semicirculatim posse eos curvare, ut insectum cui assiguntur sugendo se alant. Interdum illis adeo cooperiuntur Coleoptera, ut elytra seu alarum tegumenta non possint conjungi, ipsique artus impediti ægre moveantur.

Acari Vegetantis superficies superior est glabra et convexa; inferior sere complanata. Clarius tamen patebit similitudo insectum ipsum depictum inspicienti, quam ex descriptione colligi possit. Figuræ duæ superiores acarum anticum et possicum repræsentant. Figura autem magna ostendit plures, sex nempe, elytro adhærentes illius scarabæi quem varietatem esse simetarii ante diximus. Alia maxillas, cæterasque capitis partes mire conformatas, multo tamen auctius, monstrat.


Magnum horum me memini numerum Onisco a Linnæo Armadillo nominato, seu millepedæ officinarum, ut dicitur, annexum semel conspexisse. Ut verbo dicam, acarum vegetantem nihil aliud esse suspenter larvam acari coleoptratorum Linnæi, qui super scarabæos sere omni tempore possit inveniri, physicisque optime innotescit. Notandum præterea acarum vegetantem non in genere acari apud Linnæum reperiri, sed in appendice animalium ad siném voluminis tertii Systematis Naturæ.

Celeberrimus

Celeberrimus De Geer in Actis Stockholmienfibus anni millesimi septingentesimi sexagesimi octavi parvulum hoc animalculum descripsit et delineavit, vix tamen satis eleganter aut accurate. In Historia Insectorum iterum illud descripsit idem auctor et depinxit.







London Luttished April Plager by F. P. Nodder & P. Nis Brewer Sweet

#### THE

# VEGETATING MITE.

#### GENERIC CHARACTER.

Eight Legs.

Two Eyes, fituated on the fides of the head. Two Tentacula, jointed, and shaped like feet.

SPECIFIC CHARACTER.

CRUSTACEOUS REDDISH or YELLOW-ISH-BROWN ACARUS, convex above, flat beneath.

That curious little infect, the acarus vegetans of LINNÆUS, or the vegetating mite, being, on account of its minuteness, less attended to than many others of its genus, it cannot be unacceptable to the admirers of Natural History to see it accurately figured and described. It is generally to be found on the bodies of other infects; and particularly on those belonging to the beetle tribe, or such as in the Linnæan language, belong to the class Coleoptera.

The infect on which I have most commonly observed it, is the Hister Unicolor Lin: which is a smallish coleopterous infect, frequently seen in gardens in the early part of the spring, and which at that season seldom fails to have some of the vegetating mites either


on its body or wing cases. I have also observed it in great plenty on the scarabæus stercorarius Lin: or common black beetle, as well as on a small brown variety of the scarabæus simetarius Lin: which is almost always to be found in cow-dung in the months of April and May.

The manner in which these minute insects adhere is very curious; for each is affixed by a flexible stalk or pedicle to the wing-shells and other parts of the insect they insest; so that unless their nutriment is absorbed by the stalk which supports them, (and which, in this case, must be tubular,) they must have the power of bending the stalk in a semicircular direction at particular times, so as to enable them to adhere occasionally to the insect for the purpose of seeding. They are sometimes so numerous as to prevent the beetles on which they grow, from closing their wing-sheaths; and adhere to their limbs in such a manner as greatly to impede their motions.

The upper furface is smooth and convex; the lower surface slat; but the sigures annexed will convey a clearer idea of the several particulars of their appearance than any description. The two upper sigures represent the fore and back view of the insect. The large groupe represents six of them adhering to the elytron or wing-sheath of the brown variety abovementioned of the scarabæus simetarius. The other sigure shews the singular appearance of the jaws and other parts at the head of the insect, magnissed in a much greater degree.

I once observed these insects in great plenty on the oniscus armadillo, or common officinal Millepede. Upon the whole, I am strongly inclined to believe the vegetating mite to be nothing more than the larva or young of the acarus coleoptratorum, or common beetle-acarus, which insects those insects at most seasons of the year, and which is well known to every observer of insects. I should observe that the acarus vegetans does not occur under the genus acarus in the Systema Naturæ of Linnæus, but is mentioned in the appendix animalium at the end of the third volume of that work.

DE GEER in the Acta Stockholmiensia for the year 1768 has described and figured this insect; but his figures cannot be commended either for their elegance or accuracy. He has also figured and described it in his History of Insects.



# LACERTA PLATURA,

#### CHARACTER GENERICUS.

Corpus tetrapodum, caudatum, nudum.

Lin. Syst. Nat. p. 359.

#### CHARACTER SPECIFICUS.

LACERTA cauda depresso-plana lanceolata, margine subaculeato, corpore griseo-susception feabro.

Ungues quasi duplicati: lingua brevis, lata, integra, seu non forficata; apice autem leniter emarginato.

Lacertam Platuram, quam nuperrime detectam generavit Nova Hollandia, infigniter distinguit cauda depressa et complanata, margine tenuissimo, sensim in acutum apicem decrescens. Rarissima est in genere hæc caudæ planities, nec accidit nisi duabus vel tribus speciebus. Uncias quatuor cum semisse paulum superat longitudine Platura. Caput magnum pro corpore. Tota superior superficies tuberculis parvis exasperatur, quæ in aliquibus partibus, versus occiput præcipue et caudam, in mucrones acuminatos extenduntur. Inserior superficies pallida est, seu albescens.

#### THE

# BROAD-TAILED LIZARD.

**◇**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*

GENERIC CHARACTER.


Body four-footed, tailed, naked.

SPECIFIC CHARACTER.

LIZARD with a depressed lanceolate tail, almost spiny on the margin; the body of a dusky grey color, and rough.

The claws appear as if double; the tongue is short and broad, not forked, but slightly emarginated at the tip.

The Lacerta Platura or broad-tailed Lizard, a species very lately discovered, and which is a native of New Holland, is strikingly distinguished by the uncommon form of its tail, which is of a depressed or slattened shape, with very thin edges, and gradually tapers to a sharp extremity. This depressed form of the tail is extremely rare in lizards, there being scarcely more than two or three other species in which a similar structure takes place. This lizard is somewhat more than sour inches and a half in length; the head is large in proportion; and the whole upper surface of the animal is beset with small tubercles, which in some parts, especially towards the back of the head, and about the tail, are lengthened into sharpened points. The lower surface is of a pale color, or nearly white.





## TROCHILUS COLUBRIS.

\*\*\*\*\*\*\*\*

#### CHARACTER GENERICUS.

Rostrum subulato-filisorme, apice tubulato, capite longius: Mandibula fuperior vaginans inferiorem.

Lingua filiformis, filis duobus coalitis tubulofa. Pedes ambulatorii.

Lin. Syft. Nat. p. 189.

# HARACTER SPECIFICUS, &c.

TROCHILUS rectirostris viridi-aureus, rectricibus nigris, lateralibus tribus ferrugineis apice albis, gula flammea.

Lin. Syst. Nat. p. 191.

MELLIVORA gula rubra.

Edw. Av. t. 36.

MELLIVORA Avis Carolinensis.

Catesb. Carol. 1. p. 65.

Coloribus nitidis adeo gemmeisque exornatur minutum Trochili genus, ut si cum his comparantur, non modo fordeant prorfus et obscurentur avium plurium regiones calidiores incolentium venustissimi colores, fed et aurum ipsum pyropusque longe superentur. Nequeunt igitur ulla arte splendidissimi hi colores ad


Gg

vivum

vivum exprimi, quæ tum fane magnum præstitit, si insigne harum avium decus imitando adumbraverit. Regulus cristatus Europæus avicula omnium minima diu habebatur, et stupebant prorsus nostrates physici, cum aves Americanas vel ipsis insectis Europeis minores conspicerent; inter quas est Trochilus vix scarabæo vulgari major. Cum tamen minuta hæc species cæteris cedat colorum splendoribus, satius duxi aliam eamque elegantiorem exprimere.

Formofissima hæc avicula plurimas Americæ partes, Carolinam præcipue tam Septentrionalem" quam Auftralem inhabitat. Rostrum illi nigrum est. Corporis fuperiora splendent aureo-viridi, et mutabili colore. Gulam decorant plumæ coccineæ nitidissimæ, quæ pro lucis ratione in aureum fulgentissimum transeunt. Pectus et abdomen albicant. Alarum remigeo fuscopurpurei. Feminæ superficies inferior albicat, carens omnino rubore illo eximio quo mas infignitur. Miranda est huic aviculæ pennarum velocitas. volandum cibum capit phalænæ more, linguam nempe tubulatam in fundum florum immergens et mel inde exugens. Pugnacissimo est ingenio, nec raro de eodem flore acerrime dimicatur. In domus frequenter involat Trochilus, cumque, more infectorum Europæorum, tres quatuorve gyros egerit, dicto citius e fenestris essugit. Ubi denfissima est arborum coma nidificat, plerumque inter furcas ramulorum. Diametros interna nido est unciæ, profunditasque unciæ dimidiatæ: tomento, seu lanugine plantarum intus struitur, extrinsecus musco et lichene stipatur. Ova tantum duo deponit hæc avis, alba, magnitudine piforum. Dum incubat





- Somber , Saltwood . They of My Stilled adder of 3. A is Browner . Il

#### THE

# JACULATOR,

O R

# SHOOTING FISH.

**\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*** 

#### GENERIC CHARACTER.

Teeth (generally) fetaceous, fmall, and very numerous.

Body commonly fasciated transversely. Dorsal and anal Fins sleshy and scaly toward the base.

#### SPECIFIC CHARACTER.

WHITISH CHÆTODON, with nine spines and an ocellated spot on the dorsal fin, and a lengthened subcylindric snout.

The fish here represented in its natural fize, is a native of India, where it inhabits rivers, especially where the water, from its nearness to the sea, is a little brack-ish. It is remarkable for the very extraordinary manner in which it takes its prey. It principally feeds on slies and other insects which hover over the waters. When it observes one of these in its passage, it shoots against it from its tubular shout a drop of water, with so much force, and with so sure an aim as generally to lay it dead on the surface. In the same manner it shoots

shoots at such insects as happen to be sitting on the tops of plants and other projecting objects near the banks. In shooting at a sitting insect it commonly approaches to the distance of from six to sour feet before it explodes the water.

When kept in a state of confinement in a large bafon of water, these sish afford a high entertainment, by their dexterity in this exercise. If a sly or other insect be fastened on the edge of the vessel, the sish instantly perceive it, and, as if contending for the prize, continue to shoot at it, with such admirable skill as scarce ever to miss their mark.

The color of this fish is whitish, or very pale brown, with (commonly) four or five blackish fasciæ running across the body. The eyes are very large. On the lower part of the dorsal fin is a very remarkable large black ocellated spot, surrounded by a circle of a lighter color. The snout is sinely calculated for the power of exploding water from it. The trivial name of Rostratus, which Linnæus applied to this sish, being not sufficiently distinctive of the species, but applying with perhaps equal propriety to some others of the genus, I have presumed to substitute that of Enceladus.

I think it necessary to observe that it is not improbable that the faculty of darting or shooting water from the snout may not be absolutely peculiar to the fish above-described, but may perhaps take place in two or three other species.

# PAPILIO ACHILLES.

\*\*\*\*\* \*\*\*\*\*\*\*\*\*\*\*\*\*\*\*

#### CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavatocapitatæ.

Alæ (sedentis) erectæ sursumque conniventes (volatu diurno).

Lin. Syst. Nat. p. 744.

#### CHARACTER SPECIFICUS.

PAPILIO alis dentatis : fupra nigris fafcia cærulea ; fubtus fufcis : ocellis tribus quinifve.

Lin Syst. Nat. p. 752.


Eq: Achiv.

Clerk. ic. t. 24. fig. 2. Mer. Sur. t. 7. ?

Americam Australem incolit grande hoc insectum, venusta quadam dignitate conspicuum. Colores interdum variant. Habent nempe nonnulla specimina sasciam istam quæ splendidissime cærulea est latiorem multo quam alia. Disserunt etiam tum numero tum colore pallidæ illæ maculæ ad apices alarum superiorum sitæ, interdumque sere albæ sunt, et paululum ocellatæ, interdum subsusce obumbratæ. Supersiciei inferioris

inferioris color est fuscus, striis pallidioribus undulatus, maculisque magnis ocellatis, quarum alba sunt centra, distinctus.





# ACHILLES,

#### ORTHE

# GREAT BLUE-BANDED BUTTERFLY.

\*\*\*\*\*\*\*\*\*\*

#### GENERIC CHARACTER.

The Antennæ or Horns thickening towards the upper part, and generally terminating in a knob, or club-shaped tip.


The Wings (when fitting) erect, and meeting upwards. (Flight diurnal).

#### SPECIFIC CHARACTER.

BLACK BUTTERFLY with indented wings, croffed by a broad band of blue: the lower furface brown, with ocellated fpots.

This infect, so remarkable for its size and the grandeur of its appearance, is a native of South America. It is subject to some variety; the band of brilliant blue being wider in some specimens than in others: there is also a difference in the number as well as the color of the pale spots with which the upper wings are marked towards their tips, which in some specimens are nearly white, and of a somewhat occllated appearance; in others

others tinged with pale brown. The lower furface of this infect is brown, undulated with paler streaks, and ornamented by some large ocellated spots with white centres or pupils.





### ANAS MERIANÆ.

\*\*\*\*\*\*\*

#### CHARACTER GENERICUS.

Rostrum lamelloso-dentatum, convexum, obtusum.

Lingua ciliata, obtufa.

Lin. Syst. Nat. p. 194. Ord. Anseres.

#### CHARACTER SPECIFICUS.

ANAS cupreo-ferrugineo cyaneo viridique varia, fubtus alba, capite collo postico remigumque secundariorum apicibus carunculatis.

Ab exemplari, quod cum aliis multis Surinamensibus a lectissima Merian eximie depictis, in Museo Britannico affervatur, pulcherrimam hanc avem imitati sumus.

Omnes quibus illam intueri datum est semper torsit species incognita. Ad Linnæi Anatis moschatæ similitudinem, seu, ut vulgo dicitur Moscoviticæ videtur accedere; caput tamen pluribus tuberculis aspergitur, per collum etiam posticum longe decurrentibus; quodque magis mirandum, in apicibus remigum secundariorum magna horum exuberat luxuria, sasciam duplicem caruncularum per mediam alam transverse H h

ducentium. Effecit res nova et inaudita ut ipfius picturæ veritas in dubium vocetur, cum in reliquo hoc genere frustra quæratur simile quidpiam, nec sane in alio aliquo tanta sit tuberculorum ubertas.

Fatendum est Garruli Boiohemici, qui Ampelis Garrulus Linnæi, remiges fecundarios appendicibus complanatis corneis ruberrimifque terminari; quod et in Gallo gallinaceo, si juxta naturam ipsam vixerit. minus tamen perspicue, cernitur; in extremitatibus nempe pennarum quæ in collo funt. Annon igitur in hac anatis specie remigibus secundariis revera appendantur tubercula; feu habeat avis certis temporibus carunculas laterales, quæ per ipfas alarum pennas interdum protrudantur; an denique ipsa Merian, ut phyficorum ingenium exerceret, avemque formofiorem redderet, a veritate semel sciens volensque deflexerit; hæc omnia futuris indagatoribus relinquamus necesse est. Certum interim est hodiernis physicis plane ignotam esse hanc avem, illamque ex antiquioribus neminem aut descripsisse aut depinxisse. Magnitudine Anatis communis feu Boschadis a Meriana depingitur.





#### THE

## MERIAN DUCK.

\*\*\*\*\*\*\*\*

#### GENERIC CHARACTER.

Bill broad and flattened; the edges marked with fharp lamellæ.

Tongue broad and ciliated at the edges.

#### SPECIFIC CHARACTER.

DUCK varied with copper-ferruginous, deep blue, and green: white beneath; with the head, back of neck, and tips of fecondary wing-feathers carunculated.

The very extraordinary and beautiful bird here figured is accurately copied from a most elegant drawing by the celebrated Madam Merian, and is one of the birds of Surinam which occur in the representations by that lady now preserved in the British Museum.

This bird has uniformly puzzled every ornithologist who has viewed it, to determine its real species. It is evident at first sight that it bears a strong affinity to the Anas moschata of Linnæus, which is commonly called the Moscovy Duck; but is much more tuber-culated about the head, and even a very considerable distance down the back of the neck; and what is to the last degree singular, the same sort of tubercles appear

pear

It is true that in the bird called the Ampelis Garrulus, or Bohemian Chatterer, the tips of the secondary wing-feathers are terminated by flat callous or horny appendages of a bright red color; and an appearance in some degree analogous to this takes place also in the tips of the neck-feathers of the common cock in a state of Nature. Whether therefore in this species of duck the tips of the fecondaries be really furnished with tuberculated appendages; or whether the bird, at particular feafons may not be furnished with lateral caruncles, which may occasionally protrude between the feathers of the wings as represented by Madam MERIAN; or lastly, whether that ingenious lady may not in this inflance have departed a little from her general accuracy, and have given what she might have confidered as an additional ornament, must be left to future enquiries to determine: certain it is that the bird is unknown to modern ornithologists, and is neither figured or described by any other natural historian. The fize of Madam Merian's figure is nearly that of a common duck.

## HEDYSARUM GYRANS.

\*\*\*\*\*

CHARACTER GENERICUS.

Corollæ carina transverse obtusa. Legumen articulis monospermis.

Lin. Syst. Nat. p. 493. Diadelph: Decandr:

CHARACTER SPECIFICUS.

HEDYSARUM foliis ternatis.

Lin. Suppl. Plantar. 1781. p. 332.

Cum paucis abhinc annis ab interiori Bengala ad nos pervenerit hæc planta, perculit omnino Europæos physicos mira illius atque antehac incognita qualitas; motus nempe perpetuus et spontaneus, causis externis nec impeditus nec acceleratus, Mimosæ aliarumque nonnullarum more negans irritari. Efficitur hic motus alternatim convenientibus et recedentibus duobus parvulis foliis utrinque ad pediculos sitis, quæ per totum sere diem leni hoc exercitio agitantur.


Apud nos tamen non femper expectationi respondet Hedysarum Gyrans, frigus sentiens torpidum, aeremque pigrum et a nativo longe diversum, manensque meridiem, languido et imbecillo conatu vim insitam ægre evocat: clarissimum argumentum mitiores zephyrorum auras, mollemque cœli Indici temperiem hypocaustorum vaporibus male permutari. In Gangetica terra facili-

facillimus et liberrimus est hujus plantæ motus, quam facram habent incolæ, cæterisque longe digniorem.

Spontaneo huic motui nihil adhuc in vegetabilibus fimile repertum est: inter plurima est quæ oculis, non intellectu cernuntur, quæque non possumus non admirari, ignorantiam fatentes.

In Europa ad altitudinem duorum vel trium pedum plerumque crescit, foliaque gerit læte viridia, quorum media pars magis est glauca; floresque rubentes subcæruleo tinctos, et interdum flavescentes.





# ANIMATED HEDYSARUM,

OR THE

#### MOVING PLANT.

**\*\*\*** 

GENERIC CHARACTER.

Keel of the *Corolla* transversely obtuse. *Legume* with single-seeded joints.

SPECIFIC CHARACTER.

HEDYSARUM with ternate leaves.

The plant of which a figure is here given was fome years backwards transmitted to us from the interior parts of Bengal, and struck the European Naturalists with astonishment at a phænomenon till then unknown amongst vegetables; viz. a constant and voluntary motion; unconnected with any peculiar irritability, as in the sensitive plants and some others, and neither accelerated or retarded by external causes. The motion consists in the alternate meeting and receding of the two small appendages or leasets situated on each side the footstalks, and which are engaged in this gentle exercise during the greatest part of the day.

In our own country, however, it is not always that the specimens of this curious plant, in an air so different from that of their native regions, exhibit themselves to advantage: they frequently feel the benumbing effects fects of an unfavorable climate, and only make a faint and feeble attempt towards the middle of the day, at exerting their extraordinary faculty; a convincing proof how much the artificial heat of a northern flove is inferior to the genial warmth and balmy foftness of its native Indian atmosphere! It is there that it exerts its wonderful motions with unrestrained freedom, and is regarded as a facred plant, possessed of powers superior to the common race of vegetables.

This voluntary motion is not analogous to any other yet observed in plants: it is one of those numerous phænomena which we are obliged to view without understanding, and to admire without being able to explain.

The specimens raised in Europe generally rise to about the height of two or three seet: the leaves are of a bright green with the middle part of a more glaucous appearance than the rest: the flowers are of a pale red, slightly tinged with blueish, and sometimes yellowish.

#### SERTULARIA SETACEA.

\*\*\*\*\*\*\*\*\*\*\*

CHARACTER GENERICUS.

Flores Hydræ.

Stirps radicata, fibrofa, nuda, articulata: articulis unifloris.

Lin. Syst. Nat. p. 1306.

CHARACTER SPECIFICUS, &c.

SERTULARIA fimplex pinnata, pinnis alternis fubincurvatis, denticulis obfoletis remotiffimis fecundis, ovariis oblongo-tubulatis axillaribus.

Ellis. Zooph. p. 47.

SERTULARIA PINNATA. B.

Lin. Syst. Nat. p. 1312.

Genus Sertulariæ, exemplum admirabile exstat animalis sub specie vegetabilis. Tam apprime enim plantas repræsentant pleræque Sertulariæ species, ut pene ab omnibus pro muscis marinis habitæ, itaque ab historiæ naturalis studiosis descriptæ sint. Solertissimus autem nuper desunctus Ellistus plane probavit hæc simulata vegetabilia veras esse hydras, seu polypos marinos; ab hydris aquarum dulcium in hoc differentes, quod habeant corneam velut thecam corporum ramu-

los


los cingentem, defendentemque ab omnibus injuriis quas in elemento tam turbido et feroci nudæ experirentur.


Sertulariæ rupibus, conchis aliisque, stolonibus, seu radicibus velut repentibus assiguntur; et hoc modo plantis videntur simillimæ. Corpuscula illa ovata, quæ in axillis ramulorum conspiciuntur, pro ovariis ab Ellisio habentur, credunturque continere ova, et interdum etiam sætus plane formatos. Species Sertulariæ numerosissimæ exstant, quarum plures sunt vulgatissimæ.

Species quam hic depingi curavimus ab Ellisio mutuati fumus, utpote structuræ qualitates peculiares probe exprimentem. Pars animalis, velut medulla, undique per corneam thecam disfusa est; et ex singulo ramulorum denticulo sub forma capitis hydræ protruditur cum suis tentaculis. Ovaria, ut vocat Ellisius, in axillis ramorum sita sunt. Sertulariæ, sicut hydræ aquarum dulcium, omnia animalcula prædantur quæ tentaculis suis sors obtulerit.

Fig. 1. Animal magnitudine naturali.

Fig. 2. Idem microfcopio auctum.





# THE SMALL SEA-BRISTLE CORALLINE.

�\*\*\*\*\*\*\*\*

GENERIC CHARACTER.

A compound Hydra or Polype ramified in the manner of a plant, included in a horny case, and affixed by its base.

SPECIFIC CHARACTER, &c.

SIMPLY PINNATED SERTULARIA, with bent alternate pinnæ furnished with very remote processes growing on one side only, and oblong axillary ovaries.

LITTLE SEA-BRISTLE CORALLINE.

Ellis Zooph. p. 47.

Ellis's Corallines, pl. 38. fig. 4.

The genus Sertularia affords a most curious proof of an animal body under the appearance of a vegetable. So very great is the similarity which most of the species bear to plants, that they have almost universally been regarded as a fort of sea-mosses, and as such have been described by most naturalists; but the late ingenious Mr. Ellis seems to have clearly proved that these apparent vegetables are no other than real marine polypes or hydras; which differ from the fresh-water polypes in being provided by Nature with a horny case

case or tube, accompanying the ramifications of their bodies, and serving to defend them from the numerous disasters to which they would otherwise be liable, if left naked in the tumultuous element in which they are destined to reside.

They adhere to rocks, shells, &c. by creeping processes, which bear the appearance of spreading roots, and thus contribute still farther to their plant-like aspect. The oval bodies which are so frequently seen seated at the bases of the lateral branches, are supposed by Mr. Ellis to be the ovaries of the animal, containing the eggs, and sometimes the completely-formed young. The species of Sertularia are extremely numerous, and are very common.

The species here figured we have copied from the works of Mr. Ellis, as it serves to shew in the clearest and most distinct manner possible, the several particulars of the supposed structure. The animal part like a medulla or pith, is every where seen through the transparent horny coat; and at every denticulation of the branches it is protruded in the form of a polype-head with its tentacula. The supposed ovaries are situated in the axillæ of the branches. The Sertulariæ, like the fresh-water polypes, prey on such small animalcules as happen to fall in the way of their tentacula.

Fig. 1. reprefents the animal in its natural fize.

Fig. 2. shews it magnified by a microscope.

#### STRUTHIO RHEA.

�\*\*\*\*\*\*\*

CHARACTER GENERICUS.

Rostrum subconicum, depressum. Nares ovatæ. Alæ ad volandum ineptæ. Pedes cursorii.


CHARACTER SPECIFICUS.

STRUTHIO pedibus tridactylis.

Ab exemplari pulcherrimo Americano, in Museum Leverianum nuperrime illato nunc primum depingitur avis exoticarum fere omnium rarissima, quamque probabile est non modo non vidisse Europæos physicos, sed pene ignorasse. Struthioni licet communi seu Africano primo intuitu fimillima fit Rhea, alarum tamen pennæ, utcunque ad volandum inutiles, non modo longe majores funt, fed et laxæ et quasi divaricatæ; et utrinque super axillas exstat fasciculus plumosus elongatus, torquem efficiens, quali propemodum distinguitur cervix Tetraonis Umbelli. Facillime vero speciem denotat, Linnæoque pro charactere specifico inserviit pedum forma, qui tridactyli sunt, digitis omnibus Kk

omnibus antrorsum spectantibus, ungulatis, et æquali fere magnitudine; cum habeat pes Struthionis communis duos digitos, quorum major tantum unguibus instruitur. Color Rheæ, in hoc saltem specimine, est terreo-suscus, subtus pallidior, pennis alarum intermediis seu interioribus albis. Cauda vix discerni potest; in uropigio tamen paulo productiores sunt pennæ quam in reliquo corpore. Crura pedesque nigricant.

Notandum est in America tantum Australi reperiri hanc speciem, et pullum forsitan suisse avem quæ in Museo Leveriano asservatur; est enim Struthione Africano dimidio minor.





#### THE

## AMERICAN OSTRICH,

GENERIC CHARACTER.

Bill depressed, rounded at the end. Wings useless for slight.

Legs naked above the knee.

SPECIFIC CHARACTER.

OSTRICH with three-toed feet.

The Struthio Rhea, or American Ostrich, one of the rarest of exotic birds, and which appears to have been hitherto almost unknown in Europe, and scarce ever seen by any European naturalist, and which most certainly has never before been figured, is here faithfully represented from a very sine specimen lately sent from America to the Leverian Museum. In its habit, or general appearance, this bird at first sight bears a near resemblance to the common or African Ostrich; but the wings, though useless for slight, are composed of seathers which greatly exceed those of the common ostrich: they are also looser, and more divaricated or K k 2

fpread than in that species; and on each fide the neck, just above the axillæ, is a fasciculus of long feathers, forming a fort of ruff, almost in the same manner as in the Tetrao Umbellus, or ruffed heathcock; but what eafily diffinguishes this species, and which LINNEUS has made use of for his specific character of the bird, is, that the feet are not didactylous, or composed of two toes, as in the common or African species, but are tridactylous, or composed of three distinct toes; all which point forwards, and are nearly alike as to proportion: they are likewife all three furnished with claws; whereas in the common offrich there are only two toes, of which the primary or large one alone is clawed. The colour of this bird (at least in the present fpecimen) is earthy-brown, paler beneath, and the intermediate or interior wing-feathers are white. There is fcarce any appearance of a tail, but the feathers on the rump are a very little longer than in other parts. The legs and feet are blackish. I should observe, that this specimen is probably not a full-grown one, and is scarcely half the fize of the common or African offrich.

## MUS STRIATUS.

CHARACTER GENERICUS.

Dentes Primores inferiores subulati.

Lin. Syft. Nat. p. 79.

CHARACTER SPECIFICUS, &c.

MUS rufus, corporis striis pluribus albo-guttatis.
MUS Orientalis.

Seba. 2. t. 21. f. 2. Pall. Glir. p. 90. No. 97.

In calidioribus ut plurimum mundi partibus, in India præcipue Orientali reperitur mus hicce parvulus et pulcherrimus. Color est rusus seu ferrugineosus subtus pallidior magisque albicans. Ornantur dorsum et latera crebris ordinibus punctorum alborum quæ formam ovatam obtinent. Crura quoque et cauda subalbida sunt. Cum ingenio sit Mus striatus miti admodum et innocuo facillime mansuescit. Magnitudine muri vulgari paululum cedit.

#### THE

## STRIPED MOUSE

**~** 

#### GENERIC CHARACTER:

Two cutting teeth in each jaw. Four toes before; five behind. Slender taper tail.

Specific Character, &c.


RUFOUS MOUSE, with the body marked by feveral rows of oval white spots.

ORIENTAL RAT.

Penn. Quadr. p. 445.

This little animal, so eminently distinguished by the beauty of its appearance, is chiefly sound in the warmer regions of the globe, and particularly in the East-Indies. Its colour is a ferruginous or reddish brown, paler, or more inclining to white beneath, and decorated on the back and sides by several rows of small oval spots of white. The tail and legs also incline to whitish. It is easily tamed, and its disposition is perfectly mild and gentle. It is somewhat less than the common mouse.

Landon , Sublished July 1999 by St. P. Salder & C. S. Brown Some .



8



## COLUBER NAJA.

CHARACTER GENERICUS.

Scuta abdominalia.
Squamæ fubcaudales.

Lin. Syst. Nat. p. 275.

CHARACTER SPECIFICUS.

COLUBER FERRUGINEO-FLAVESCENS, collo (plerumque) multum dilatato macula-que fupra conspicillata albo nigroque varia notato.

Scuta abdominalia circiter 193. Squamæ fubcaudales 60.

Inter Europæorum commoda jure potest reputari serpentum venenosorum paucitas, qui in Indiæ et Africæ vastis regionibus, nec non in Americæ late patentibus desertis denso et horrendo dominantur agmine; nonnulli in exitium humani generis nimis, heu! lethaliter armati.

Omnes virulenti ferpentes dentibus tubulatis venenum per foramen e facculo maxillari in vulnus inflantibus, inftructi funt. Sunt tamen alii illis characteres quibus aliquatenus ab innoxiis ferpentibus diftingui

tingui possunt. Notavit Dominus Gray, in Actis Anglicis plerosque venenosos serpentes caput latum, depressum, squamis minutis tectum habere; corporaque squamis carinatis, seu linea elevata notatis obducta: at e contrario, innoxiis ferpentibus, qui morfu tantum fimplici lædunt, caput plerumque effe parvum, fquamis magnis et latis adopertum, corpusque squamis lævibus, feu non carinatis. Hi autem characteres generales habendi funt, nec funt improvide et indubitanter recipiendi. Ita fe rem habere species depicta exemplum est clarissimum. Est enim hic serpens inter pestiferos fere diriffimus; et tamen externis illis specierum perniciofarum characteribus non distinguitur; fed ut innocuo serpenti caput est illi plerumque parvum, squamis magnis tectum; corpusque squamis lævibus, seu non carinatis.

Tremendum hocce animal in India Orientali invenitur. Morfus ejus non minus quam Crotali funestus est. Ab Indiæ tamen incolis interdum capitur, et telis, seu dentibus tubulatis extractis (mirum dictu!) mansuescit, et in varios modos, ad domini nutum, velut saltans, se contorquet; et sæpe pro spectaculo exhibetur. Nota, quam in collo habet singularem, diversis speciminibus plus minus vivida est. Color etiam totius corporis in diversis aliquatenus differt. In longitudinem aliquot pedum interdum crescit hic serpens.

Cum monstra hæc, regiones calidiores orbis incolentia, animo occurrunt, gratulari nobismetipsis merito possumus de nostra securitate; et lubentissime anteponamus hyemis septentrionalis incommoda, pigrosque campos

campos et diu infructuosos, perenni climatis æstati, et sloribus perpetuo ridentibus, quibus India utraque superbit: at ubi, proh dolor! viator incautus, media inter gaudia, sato subitaneo corripitur.

#### THE

### SPECTACLE SNAKE, O R

## COBRA DE CAPELLO.

GENERIC CHARACTER.

Transverse Lamellæ under the abdomen. Broad alternate Scales under the tail.


SPECIFIC CHARACTER.

FERRÜGINOUS-YELLOW SNAKE, with the neck (generally) much dilated, and marked above by a spectacle-shaped spot of black and white.

The abdominal plates are about 193. The subcaudal scales 60.

In Europe the noxious part of the ferpent tribe is happily confined to a very few poisonous species; but the vast regions of India and Africa, and the extensive wilds of America, are infested by a variety of thefe dreadful reptiles; fome of which are but too well provided with the fatal power of destroying mankind by their bite.

Such





Such ferpents as are of a venomous nature, are furnished with tubular fangs, or teeth, through which their poison, which is contained in refervoirs at the roots of the fangs, is injected into the wound. Besides the fangs, there are in general some external characters which may in fome measure ferve to diffinguish the poisonous ferpents from the innoxious ones: thus it has been observed by the ingenious Dr. GRAY, in the Philosophical Transactions, that the generality of poisonous ferpents have a broad, depressed head, covered with small scales; and that their bodies are commonly covered with carinated scales, i. e. scales which have a prominent line upon them. On the contrary, the innoxious ferpents, or fuch as can merely inflict a fimple wound, have generally a fmallish head, covered with large broad scales, and the body covered with smooth scales, or not carinated: but these are characters which admit of exceptions, and are merely to be regarded in a general view. The species here figured is a striking instance of this, and (as Dr. GRAY observes) is in every respect a complete exception to what has been faid respecting the distinction between venomous and innoxious ferpents; for though this is one of the most terrible of the whole tribe, yet it is not distinguished by those external marks of malignity which commonly characterize the poisonous species: on the contrary, it agrees with the innoxious ferpents in having a smallish head, covered with very large fcales; while the body is covered with fmooth fcales. This formidable animal is a native of the East-Indies. Its bite is not less destructive than that of the Rattle-Snake

Snake. It is faid however to be fometimes caught by the Indians, and after having its fangs drawn, to be in fome degree tamed, fo as to be taught to throw itself into various fantastic attitudes, as if daneing; and in this condition is often exhibited in India as a shew. The very singular mark on the neck is much more apparent in some specimens than in others. The colour also of the whole body is deeper or lighter in different specimens. It grows to the length of some feet.

Reflecting on these horrid natives of the hotter regions of the globe, we may congratulate ourselves on our own happy state of security, and may well be willing to prefer the rigours of a northern winter, with a temporary loss of vegetation, to the continued warmth of climate, and ever-blooming verdure, which distinguish the beautiful regions of both the Indies; where alas! the ineautious traveller may meet with sudden sate in the midst of the most enchanting scenes which nature can display.

#### INDEX.

## 42. ACARUS autumnalis.

- 64. Acarus vegetans.
- 60. Anas Merianæ.
- 51. Boa Constrictor.
- 58. 59. Bradypus urfinus.
- 41. Buceros Rhinoceros.
- 57. Chætodon armatus.
- 67. Chætodon Enceladus.
- 74. Coluber Naja.
- 40. Dionæa Muscipula.
- 62. Gryllus migratorius.
- 70. Hedyfarum gyrans.
- 46. Julus maximus.
- 45. Lacerta Salamandra.
- 65. Lacerta platura.
- 56. Loxia jugularis.
- 43. Mantis Gigas.
- 38. Mus faliens.
- 73. Mus striatus.
- 63. Muscicapa porphyrobroncha
- 49. Papilio Menelaus.
- 68. Papilio Achilles.
- 39. Pennatula phosphorea.
- 60. Petaurus australis.
- 44. Picus minimus.
- 53. Plittacus gloriofus.
- 50. Pfittacus magnificus.
- 48. Simia Sphinx?
- 71. Sertularia fetacea.
- 61. Siren lacertina.
- 54. Stapelia hirfuta.
- 47. Strix Nyctea, var.
- 72. Struthio Rhea.
- 55. Sorex bicolor.
- 66. Trochilus Colubris.
- 52. Voluta araufiaca.

#### INDEX.

- PL. CARUS autumnal.
  - 48. Baboon variegated.
  - 51. Boa great.
  - 49. Butterfly filver-blue.
  - 68. Butterfly great blue-banded.
  - 57. Chætodon long-fpined.
  - 50. Cockatoo magnificent.
  - 71. Coralline sea-bristle.


  - 63. Fly-catcher purple-throated
  - 56. Grofsbeak red-throated.

  - 66. Humming-bird red-throated
  - 70. Hedyfarum animated.

  - 65. Lizard broad-tailed.
- 68. Butterfly great blue57. Chætodon long-spin
  50. Cockatoo magnifico
  71. Coralline sea-bristle
  69. Duck Merian.
  40. Fly-trap Venus's.
  63. Fly-catcher purple56. Grossbeak red-thro
  41. Horn-bill great.
  66. Humming-bird red70. Hedyfarum animate
  67. Jaculator.
  46. Julus great.
  38. Jerboa Siberian.
  65. Lizard broad-tailed
  62. Locust wandering.
  43. Mantis giant.
  - 43. Mantis giant.
  - 64. Mite vegetating.
- 64. Mite vegetating.
  73. Mouse striped.
  47. Owl snowy, var.
  72. Ostrich American.
  39. Pennatula phosphoric.
  53. Parrot Pennantian.
  60. Petaurus southern.
  61. Siren.
  74. Snake spectacle.
  54. Stapelia fetid.
  58. 59. Sloth ursine.
  55. Shrew water.
  45. Salamander.
  52. Volute orange-striped.
  44. Woodpecker minute.














500-9 500-9



